

# Chapter: Indefinite Integrals

Que:  $f(x) = \begin{vmatrix} x & x^2 & x^3 \\ 1 & 2x & 3x^2 \\ 0 & 2 & 6x \end{vmatrix}$  find  $f'(x)$

$$f'(x) = \begin{vmatrix} 1 & 2x & 3x^2 \\ 1 & 2x & 3x^2 \\ 0 & 2 & 6x \end{vmatrix} + \begin{vmatrix} x & x^2 & x^3 \\ 0 & 2 & 6x \\ 0 & 2 & 6x \end{vmatrix} + \begin{vmatrix} x & x^2 & x^3 \\ 1 & 2x & 3x^2 \\ 0 & 0 & 6 \end{vmatrix}$$


## \* Indefinite integral

It is a ~~reverse~~ <sup>reverse</sup> process of diff.

$$\frac{d}{dx} x^2 = 2x$$

$$\int 2x dx = x^2 + d \quad \text{Constant}$$

family of curve



$$\therefore \frac{d}{dx} x^2 = 2x$$

$$\int 2x dx = x^2 + d \quad \begin{array}{l} \text{integrated} \\ \text{anti derivative of } 2x \end{array}$$

Here  $x^2$  is called Anti Derivative of  $2x$ .

$$\frac{d}{dx} (f(x) + c) = f'(x)$$

$$\int f'(x) dx = f(x) + C.$$

\* Every cont. fn is integrable but some of them are not expressible.

$$\left\{ \int \sin \frac{1}{x} dx = \text{Integrable but not expressible.} \right\}$$

$$\int f'(x) dx = f(x) + C$$

diff. rule

$$f(\text{cont. fn}) =$$

$$f(x) = f'(x)$$

\* if  $f(x)$  is cont. then  $F(x)$  is differentiable.

$$* \int x^{-1/3} dx = \frac{x^{-1/3+1}}{-1/3+1}$$

$$f(x) = \frac{1}{x^{1/3}}$$

$$= \frac{3x^{2/3}}{3} = \frac{3}{8} x^{2/3}$$

\*  $f(x) = 0$  discontin. at  $x=0$

but its integration  $f(x) = \frac{3}{8} x^{2/3}$  is cont. at  $x=0$

~~that~~



Que: 1)  $\int e^{\ln x} dx$

Ans:  $\int e^{\ln x} dx$

$$\int \sqrt{x} dx = \int x^{1/2} dx = \frac{x^{1/2+1}}{1/2+1}$$

(2)  $\int e^{-\ln x^2} dx$

Ans:  $\int e^{\ln(x^2)^{-1}} dx$

$$= \int (x^2)^{-1} dx$$

$$= \int x^{-2} dx = \frac{x^{-2+1}}{-2+1}$$

(3)  $\int \ln\left(\frac{1}{e^x}\right) dx$

Ans:  $\int \ln\left(\frac{1}{e^x}\right) dx = \int \ln e^{-x} = \int -x(\ln e) dx =$

$$= \int -x dx = -\frac{x^2}{2} + C$$

Que:  $\int \frac{(1+x)^3}{\sqrt{x}} dx$

Ans:  $\frac{1+x^3+3x^2+3x}{\sqrt{x}} = (x^{-1/2} + x^{5/2} + 3x^{3/2} + 3x^{1/2}) dx$

$$= \frac{x^{1/2}}{1/2} + \frac{x^{7/2}}{7/2} + 3 \cdot \frac{x^{3/2}}{3/2} + \frac{3x^{3/2}}{3/2} + C$$

$$2) \int e^{\ln 2 + \ln x} dx$$

$$\int e^{\ln 2x} dx = \int 2x dx = x^2 + C$$

$$3) \int e^{3 \ln x} dx$$

$$\int e^{3 \ln x} dx = e^{\ln x^3} = \int x^3 dx = \frac{x^4}{4} + C$$

## \* Second formula:

$$1. \int \frac{dx}{x} = \ln|x| + C$$

$$\text{or } \int \frac{dx}{x} = \ln|x| + C$$

$$2. \int \frac{dx}{ax+b} = \frac{\ln|ax+b|}{a} + C$$

$$x > 0$$

$$\text{Que: } \int \frac{dx}{2x-1}$$

$$(2) \int \frac{dx}{3-4x}$$

$$= \frac{\ln|2x-1|}{2} + C$$

$$= \frac{\ln|3-4x|}{-4} + C$$

$$(3) \int \frac{\sqrt{x^4 + x^{-4} + 2}}{x^3} dx$$

$$= \frac{\ln(x^3)}{\sqrt{x^4 + x^{-4} + 2}}$$

$$\int \frac{x^4 + \frac{1}{x^4} + 2}{x^4}$$

$$= \int \left(x^2 + \frac{1}{x^2}\right)^2 = \int \frac{x^2 + \frac{1}{x^2}}{x^3} dx$$

$$= \int \left(\frac{1}{x} + \frac{1}{x^5}\right) dx = \ln|x| + \frac{x^{-5+1}}{-5+1} + C$$

Que 0  $\int \frac{2x+3}{x^2+3x-10} dx$

$$\int \frac{(x+5)(x-2)}{(x+5)(x-2)} dx = \int \frac{1}{x-2} dx + \int \frac{dx}{x+5}$$

$$= \ln|x-2| + \ln|x+5| + C$$

Que 1  $\int \frac{x}{x^2+2x+1} dx$       (2)  $\int \frac{dx}{x^2-5x+6}$

$$\int \frac{x}{x^2+n+n+1} = \int \frac{x}{x(x+1)(x+1)}$$

$$\int \frac{(x+1) - (x+1)}{(x+1)(x+1)} = \int \frac{1}{x+1} dx + \int \frac{-1}{x+1} dx = \ln|x+1| - \ln|x+1| + C$$



$$\text{Ans: } \int \frac{(n+1)-1}{(n+1)^2} = \int \frac{dn}{n+1} - \int \frac{dn}{(n+1)^2} = \ln(n+1) \\ = \frac{(n+1)^{-2+1}}{-2+1} + C$$

$$(2) \int \frac{1}{(n-2)(n-3)} = \int \frac{(n-2)(n-3)}{(n-2)(n-3)} = \int \frac{1}{n-3} dn - \\ - \int \frac{dn}{n-2} = \ln(n-3) - \ln(n-2) + C$$

$$\text{Que. } \int \frac{dn}{\sqrt{n+1} + \sqrt{n-1}}$$

$$\frac{\sqrt{n+1} - \sqrt{n-1}}{\sqrt{n+1} - \sqrt{n-1}} =$$

$$\int \frac{dn}{\sqrt{n+1} + \sqrt{n-1}} \times \left( \frac{\sqrt{n+1} - \sqrt{n-1}}{\sqrt{n+1} - \sqrt{n-1}} \right)$$

$$\frac{1}{2} \int (\sqrt{n+1} - \sqrt{n-1}) dn = \frac{1}{2} \left[ \frac{(n+1)^{3/2}}{\frac{3}{2}} - \frac{(n-1)^{3/2}}{\frac{3}{2}} \right] + C$$

\* Third formula

$$1. \int e^n dn = e^n + C$$

$$2. \int e^{(an+b)} dn = \frac{e^{(an+b)}}{a} + C$$

$$3) \int a^n dx = \frac{a^n}{\ln a} \quad \boxed{a > 0}$$

Que:  $\int e^{2x-3} dx$

$$= \frac{e^{2x-3}}{2} + C$$

(2)  $\int 2^{3x+4} dx$

$$= \frac{2^{3x+4}}{3 \ln 2} + C$$

$\frac{2^{3x+4}}{3 \ln 2}$  ans

(3)  $\int 3^{-x} dx = \frac{3^{-x}}{-\ln 3} + C$

Que:  $\int \frac{2^{n+1} - 5^{n-1}}{10^n} = 10^n = (2 \cdot 5)^n = 2^n \cdot 5^n$

$$= \frac{2^n \cdot 2 - \frac{5}{5}}{2^n \cdot 5^n} = 2 \int 5^{-n} dx - \frac{1}{5} \int 2^{-n} = 2 \int \frac{5^{-n}}{-\ln 5} - \frac{1}{5} \frac{2^{-n}}{-\ln 2} + C$$

2)  $\int 2^n \cdot e^n dx$

$$a^n \cdot b^n = (ab)^n = \int 2^n \cdot e^n dx = \int (2e)^n dx = \frac{(2e)^n}{\ln(2e)} + C$$

3)  $\int (2^n + 3^n)^2 dx = (2^n)^2 + (3^n)^2 + 2 \cdot 2^n \cdot 3^n = \int (2^{2n} + 3^{2n} + 2 \cdot 6^n) dx$



$$= \frac{2^{2x}}{2 \ln 2} + \frac{3^{2x}}{2 \ln 3} + \frac{2 \cdot 6^x}{\ln 6} + C$$

Ques: 1)  $\int \frac{e^{3x} + e^{5x}}{e^x + e^{-x}} dx$

$$= \frac{e^{3x} (1 + e^{2x})}{\frac{e^{2x} + 1}{e^x}} = \int e^{4x} dx = \frac{e^{4x}}{4} + C$$

2)  $\int a^{mx} \cdot b^{nx} dx$

$$= \int (a^m \cdot b^n)^x = \frac{(a^m \cdot b^n)^x}{\ln(a^m \cdot b^n)} + C$$

3)  $\int 2^{\ln x} dx$

$$\left[ \begin{array}{l} \log_b e \\ a^x \end{array} \right]$$

$$= \int x \ln 2 dx = \frac{x(\ln 2 + 1)}{(\ln 2 + 1)} + C$$

## \* Fourth formula

$$1. \int \sin x \, dx = -\cos x$$

$$2. \int \sin(ax+b) \, dx = \frac{-\cos(ax+b)}{a}$$

$$3. \int \cos x \, dx = \sin x$$

$$4. \int \cos(ax+b) \, dx = \frac{\sin(ax+b)}{a} + C$$

Ques

$$1. \int \sin(2x-3) \, dx = \frac{-\cos(2x-3)}{2}$$

$$2. \int \cos(3x+4) \, dx = \frac{\sin(3x+4)}{3}$$

$$3. \int \cos^2 x \, dx = \sin^2 x + C \quad \times$$

This is wrong

$$= \int \cos x \cdot d(\cos x) = \frac{(\cos x)^2}{2} + C \quad \checkmark$$
$$\rightarrow \int \left( \frac{1 + \cos 2x}{2} \right) dx = \frac{x}{2} + \frac{1}{2} \frac{\sin 2x}{2} + C \quad \checkmark$$

$$\begin{aligned}\cos 2n &= 2\cos^2 n - 1 \\ &= 1 - 2\sin^2 n\end{aligned}$$

$$\boxed{\int \sin^4 n \, dn}$$

H.W

$$Q. \int \sin^2 n \, dn = \int \frac{1 - \cos 2n}{2} \, dn = \frac{1}{2} \left[ n - \frac{\sin 2n}{2} \right] + C$$

$$Q. \int 4 \cos^4 n \, dn$$

$$(2\cos^2 n)^2 = (1 + \cos 2n)^2 = 1 + \cos^2 2n + 2\cos 2n$$

$$= 1 + 2\cos 2n + \frac{1 + \cos 4n}{2}$$

$$= \frac{3}{2} + 2\cos 2n + \frac{1}{2} \cos 4n$$

$$\int 4 \cos^4 n = \frac{3}{2} n + \frac{2 \cdot \sin 2n}{2} + \frac{\sin 4n}{8} + C$$

$$Q. \int \sin^3 n \, dn$$

$$[\cos 3n = 4\cos^3 n - 3\cos n]$$

$$\int \frac{3\sin n - \sin 3n}{4} \, dn = \frac{3}{4} (-\cos n)$$

$$[\sin 3n = 3\sin n - 4\sin^3 n]$$

$$+ \frac{\cos 3n}{12} + C$$

$$Q. \int \cos^3 n \, dn$$

$$= \frac{-3}{4} \sin 3n - \frac{3}{4} \sin n + C$$

Ques:  $\int |1 + \sin n| dn$   $n \in (0, \pi/2)$

Ans:  $\int \frac{\sin^2 n}{2} + \frac{\cos^2 n}{2} + 2 \sin \frac{n}{2} \cos \frac{n}{2} = \int \left( \sin \frac{n}{2} + \cos \frac{n}{2} \right)^2 = \cos \frac{n}{2} + \sin \frac{n}{2}$

$= \int \left( \cos \frac{n}{2} + \sin \frac{n}{2} \right) dn = 2 \left( \frac{\sin n}{2} - \frac{\cos n}{2} \right) + C$

2)  $\int \cos 2n \cos 3n dn$

$\frac{1}{2} \int (2 \cos 5n + \cos n) dn = \frac{1}{2} \left[ \frac{\sin 5n}{5} + \sin n \right] + C$

3)  $\int \cos n^\circ dn = \frac{\sin n}{180} + C$

(n is in Radian)

$180^\circ = \pi$

$1^\circ = \frac{\pi}{180} \Rightarrow n^\circ = \frac{n\pi}{180}$

$\int \cos n^\circ dn =$

$= \int \frac{\cos \frac{n\pi}{180}}{180} dn$

$= \frac{\sin \frac{n\pi}{180}}{\frac{\pi}{180}} + C$

Ques:  $\int \frac{\cos n - \cos 2n}{1 - \cos n} dn$

$$= \frac{\cos n - 2\cos^2 n + 1}{1 - \cos n}$$

$$= \int (2\cos n + 1) dn = 2 \sin n + n + C$$

$$= \int \frac{(\cos n + \sin n)(\cos^2 n + \sin^2 n - \sin n \cos n)}{(\cos n + \sin n)} dn$$

$$= \int (1 - \frac{1}{2} \sin 2n) dn = n + \frac{\cos 2n}{2} + C$$

Ques:  $\int \frac{\cos^3 n + \sin^3 n}{\cos n + \sin n} dn$

$$= \int \frac{(\cos n + \sin n)(\cos^2 n + \sin^2 n - \sin n \cos n)}{(\cos n + \sin n)} dn$$

$$= \int (1 - \frac{1}{2} \sin 2n) dn = n + \frac{\cos 2n}{2} + C$$

3)  $\int 4 \sin n \cdot \cos^2 n dn$

$$2 \sin 2n \cdot [\cos 2n + 1] = \int (2 \sin 4n + 2 \sin 2n) dn$$

$$= \frac{\cos 4n}{2} - \cos 2n + C$$



$$Q.1 \int \frac{\cos 2x + 2 \sin^2 x}{\cos^2 x} dx$$

$$= \int \frac{1 - 2 \sin^2 x + 2 \sin^2 x}{\cos^2 x} dx = \int \sec^2 x = \tan x + C$$

$$(2) \int \tan^2 \frac{x}{4} dx$$

$$\int (\sec^2 \frac{x}{4} - 1) dx = \frac{\tan \frac{x}{4}}{1/4} - x + C$$

$$(3) \frac{1 + \cos x}{1 + \cos x} dx$$

$$= \int \frac{1 + 2 \cos^2 \frac{x}{2} - 1}{1 - 1 + 2 \sin^2 \frac{x}{2}} dx = \int \cot^2 \frac{x}{2} dx$$

$$= \int \cot^2 \frac{x}{2} dx = \int (\operatorname{cosec}^2 \frac{x}{2} - 1) dx$$

$$= \frac{-\cot x}{1/2} - x + C$$

VI formula:

$$* \int \sec x \tan x \, dx = \sec x + C$$

$$* \int \operatorname{cosec} x \cot x \, dx = -\operatorname{cosec} x + C$$

Q.1  $\int \frac{a \sin^3 x + b \cos^3 x}{\sin^2 x \cos^2 x} \, dx$

$$a \int \frac{\sin^3 x}{\sin^2 x \cos^2 x} + b \int \frac{\cos^3 x}{\sin^2 x \cos^2 x} = a \int \tan x + b \int \cot x \operatorname{cosec} x \, dx$$

$$a \int \tan x \cdot \sec x \, dx + b \int \cot x \operatorname{cosec} x \, dx \\ = a \sec x - b(\operatorname{cosec} x + C)$$

Q.2  $\int \frac{\operatorname{cosec} x + \tan^2 x + \sin^2 x}{\sin x} \, dx$

$$\int \operatorname{cosec}^2 x \, dx + \int \sin x \, dx + \int \tan x \sec x \, dx$$



## VII formula

$$* \int \frac{dx}{1+x^2} = \tan^{-1} x$$

$$* \int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

$$* \int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1} x \quad (\text{or } \frac{\pi}{2} - \cos^{-1} x)$$

$$* \int \frac{dx}{\sqrt{a^2-x^2}} = \sin^{-1} \frac{x}{a}$$

$$* \int \frac{dx}{x\sqrt{x^2-1}} = \sec^{-1} x, \quad \int \frac{dx}{x\sqrt{x^2-a^2}} = \frac{1}{a} \sec^{-1} \frac{x}{a}$$

Ex<sup>o</sup>

$$(1) \int \frac{dx}{4+x^2}$$

$$\frac{1}{4} \int \frac{dx}{1+\frac{x^2}{4}} = \frac{1}{2} \tan^{-1} \frac{x}{2}$$

$$(2) \int \frac{dx}{\sqrt{1-9x^2}}$$

$$= \frac{\sin^{-1} 3x}{3}$$

$$(3) \int \frac{dx}{1+16x^2}$$

$$= \frac{\tan^{-1} 4x}{4}$$

$$(4) \int \frac{dx}{x\sqrt{4x^2-1}}$$

$$= 2 \cdot \frac{\sec^{-1} 2x}{2}$$

$$5) \int \frac{dx}{\sqrt{16-x^2}} = \sin^{-1} \frac{x}{4}$$

Que: 1)  $\int \frac{x^2 + \cos^2 x}{x^2 + 1} \cdot \operatorname{cosec}^2 x \, dx$

$$= \frac{x^2 + \cos^2 x \cdot \operatorname{cosec}^2 x}{x^2 + 1} = \frac{x^2 + 1 - \sin^2 x \cdot \operatorname{cosec}^2 x}{x^2 + 1} \, dx$$

$$= \int \operatorname{cosec}^2 x \, dx = \int \frac{1}{1+x^2} \, dx = -\cot x - \tan^{-1} x + C$$

$$(2) \int \frac{x^2}{1+x^2} \, dx = \frac{\tan^{-1} x}{x^2}$$

deg of  $N^r \geq$  deg of  $D^r$

$$\int \left(1 - \frac{1}{1+x^2}\right) dx = x - \tan^{-1} x + C$$

$$(3) \int \frac{dx}{\sqrt{9-4x^2}} = \int \frac{dx}{\sqrt{(3)^2 - (2x)^2}} = \frac{1}{2} \frac{\sin^{-1} \frac{2x}{3}}{3} + C$$

$$(4) \int \frac{dx}{16+25x^2} = \int \frac{dx}{(4)^2 + (5x)^2} = \left(\frac{1}{4} \tan^{-1} \left(\frac{5x}{4}\right)\right) + C$$

$$* \int \frac{dn}{(n+1)(n+2)} = \int \left( \frac{1}{n+1} - \frac{1}{n+2} \right) dn =$$

$$= \ln(n+1) - \ln(n+2)$$

Ques 11

$$\int \frac{n^4}{1+n^2} dn = \frac{\tan^{-1} n}{n^4}$$

$$\frac{(n^4-1)+1}{1+n^2} = \frac{(n^2+1)(n^2-1)+1}{(n^2+1)}$$

$$= \int \left( n^2 - 1 + \frac{1}{1+n^2} \right) = \frac{n^3}{3} - n + \tan^{-1} n$$

(a)

$$\int \frac{dn}{(n^2+1)(n^2+2)} = \int \frac{1}{(n^2+1)} - \frac{1}{(n^2+2)} dn$$

$$\tan^{-1} n (\ln(n^2+1) - \ln(n^2+2))$$

$$= \tan^{-1} n - \frac{1}{\sqrt{2}} \cdot \frac{\tan^{-1} n}{\sqrt{2}} + C$$

(b)

$$\int \frac{dn}{(n-1)\sqrt{n^2-2n-3}} = \int \frac{dn}{(n-1)\sqrt{(n-3)(n+1)}}$$

$$\frac{dn}{(n-1)\sqrt{(n-3)(n+1)}} = \frac{dn}{(n-1)\sqrt{(n+1)(n-3)}}$$

$$\int \frac{dn}{(n-1)\sqrt{(n-1)^2-2^2}} = \frac{1}{2} \sec^{-1} \left( \frac{n-1}{2} \right) + C$$

Integration formula Learn

Ques:  $\int \frac{dx}{(x^2 - 4x + 4)(x^2 - 4x + 5)}$

$$\int \left( \frac{1}{x^2 - 4x + 4} - \frac{1}{x^2 - 4x + 5} \right) = \int \frac{dx}{(x-2)^2} - \int \frac{dx}{(x-2)^2 + 1}$$

$$= \frac{1}{x-2} - \tan^{-1}(x-2) + C$$

(2)  $\int \frac{dx}{(2x-7)\sqrt{(x-3)(x-4)}}$

$\begin{aligned} & x^2 - 7x + 12 \\ & x^2 - 2 \cdot \frac{7}{2}x + \left(\frac{7}{2}\right)^2 + 12 - \left(\frac{7}{2}\right)^2 \\ & (x - \frac{7}{2})^2 - \frac{1}{4} \end{aligned}$

$$\int \frac{dx}{(2x-7)\sqrt{x^2 - 7x + 12}} = \frac{1}{2} \int \frac{dx}{(x - \frac{7}{2}) \cdot \sqrt{(x - \frac{7}{2})^2 - (\frac{1}{2})^2}}$$

$$= \frac{1}{2} \cdot \frac{1}{\frac{1}{2}} \sec^{-1} \frac{(x - \frac{7}{2})}{\frac{1}{2}}$$

Ques:  $\int (x^2 + 1)^3 dx$

$$= \frac{(x^2 + 1)^4}{4} = \frac{x^7}{7} + x + \frac{3x^5}{5} + \frac{3x^3}{3} + C$$

(2)  $\int \frac{dx}{\sin^2 x \cos^2 x} = \frac{\sin^2 + \cos^2}{\sin^2 x \cos^2 x}$

$$\int \sec^2 x dx + \int \csc^2 x dx$$

$$= \tan x - \cot x + C$$

(M-2)  $\frac{4}{\sin^2 2x} dx = 4 \int \csc^2 2x dx$

$$= 4(-\cot 2x) + C$$

(3)  $\int \frac{\sin^6 x + \cos^6 x}{\sin^2 x \cos^2 x}$

(3) Ans  $\sin^6 x + \cos^6 x = 1 - 3 \sin^2 x \cos^2 x$   
 $\sin^4 x + \cos^4 x = 1 - 2 \sin^2 x \cos^2 x$

$$= \int \frac{1 - 3 \sin^2 x \cos^2 x}{\sin^2 x \cos^2 x} dx = \int \frac{dx}{\sin^2 x \cos^2 x} - \int 3 dx$$

Ques 11)  $\int \frac{(x^3+8)(x-1)}{x^2-2x+4} dx = \int \frac{(x^3+8)(x-1)}{x^2-2x+4} dx$   $a^3+b^3 = (a+b)(a^2+ab+b^2)$

$$\int \frac{(x+2)(x^2+x-2)(x-1)}{(x^2-2x+4)} dx = \int (x^2+x-2) dx$$

(2)  $\int \frac{x^3}{(x+1)} dx = \int \frac{(x^3+1)-1}{x+1} dx$


$$\frac{(x+1)(x^2-x+1)-1}{x+1} = \int (x^2-x+1) - \frac{1}{x+1} dx$$

(3)  $\int \frac{dx}{\sin^2 x \cos x}$

$$\int \frac{\sec^2 x + \cos x^2}{\sec^2 x \cos x} dx = \int (\sec x + \cot x \cos x) dx$$

21/06/17

119



Integration Virul Substitution:

If  $\int f(x) dx = \phi(x)$

then  $J = \int f(\psi(z)) \cdot \psi'(z) dz$

$\int f(k) dk$

$\frac{d\psi(z)}{dz} = \frac{dk}{dz}$

$\psi'(z) dz = dk$

$= \phi(k) + C$

$\psi'(z) = \frac{dk}{dz}$

$= \phi(\psi(z))$

$\psi'(z) dz = dk$

$2x = \frac{dk}{dx}$

$x^2 = k$

$2x dx = dk$

Ex:  $J = \int x \cos x^2 dx$

$= \frac{1}{2} \int \cos k \cdot dk$


$= \frac{1}{2} \sin k + C$

$= \frac{1}{2} \sin x^2 + C$

$2x dx = dk$

$\frac{d}{dx} x^2 = 2x$   
 $dx = \frac{dk}{2x}$

$2x dx = dk$



Formula:

1.  $\int \tan x dx = \ln(\sec x) + C$

2.  $\int \cot x dx = \ln(\sin x) + C$

3.  $\int \sec x dx = \ln(\sec x + \tan x)$  or  $\ln \tan \left( \frac{\pi}{4} + \frac{x}{2} \right)$

4.  $\int \csc x dx = \ln(\csc x - \cot x)$  or  $\ln \left( \frac{1 - \cos x}{\sin x} \right) + C$

Ques 1  $\int \frac{\tan(\ln x)}{x} dx$

$\ln x = t$   
 $\frac{1}{x} dx = dt$

Ans:  $= \int \tan t dt = \ln(\sec t) + C$   
 $= \ln(\sec(\ln x)) + C$

2)  $\int \frac{\tan(\sin^{-1} x)}{\sqrt{1-x^2}} dx$

$\sin^{-1} x = K$   
 $\frac{1}{\sqrt{1-x^2}} dx = dK$

$= \int \tan K dK$

(3)  $\int e^x \sin e^x dx$

$e^x = K$   
 $e^x dx = dK$

$= \int \sin K dK = -\cos K + C$   
 $= -\cos e^x$

$\sin K dK$   
 $-\cos K + C$

Ques 4  $\int \sec^2 x \cdot \sqrt{5+t} \cdot \tan x dx$  ( $5 + \tan x = k$ )  
 $\sec^2 x dx = dk$

$= \int \sqrt{k} dk = \frac{k^{3/2}}{3/2} + C$

$5 + \tan x = t^2$   
 $\sec^2 x dx = 2t dt$

$t = \sqrt{5 + \tan x}$

(5)  $\int \frac{x^5}{1+x^2} dx$

$= \frac{1}{6} \int \frac{d^6}{1+d^2} = \frac{1}{6} \tan^{-1} d$

$x^6 = d$   
 $6x^5 dx = dd$   
 $x^5 dx = \frac{1}{6} dd$

(6)  $\int \sec x \cdot \ln(\sec x + \tan x) dx$

$$\frac{1}{\sec x + \tan x} \cdot (\sec x + \tan x + \sec^2 x) dx = dP$$

$$\sec x dx = dP$$

$$= \int$$

Q.7)  $\frac{x^2 - \tan^{-1} x^3}{1+x^6} dx$

$$= x^3 = k$$

$$3x^2 dx = dk$$

$$x^2 dx = \frac{1}{3} dk$$

$$= \frac{\tan^{-1} x^3 = \lambda}{1+x^6} \times 3x^2 dx = d\lambda$$

$$I = \frac{1}{3} \int \lambda d\lambda$$

$$= \frac{1}{3} \frac{\lambda^2}{2}$$

(8)  $\int \frac{\int \tan x}{\sin 2x} dx = \int \frac{t}{2 \left( \frac{\sin x}{\cos x} \right) \cos^2 x} \cdot \frac{2t dt}{1+t^4}$

$$\int \frac{t}{2t^2} \cdot \sec^2 x \cdot \frac{2t dt}{1+t^4} =$$

$$= \int \frac{t}{2t^2} \cdot \frac{2t dt}{1+t^4} (1+t^4)$$

$$= \int dt = \frac{t}{\cancel{1+t^4}} = \sqrt{\tan x + 1}$$



$$Q. \frac{du}{\cos(n-1) \cos(n-2)}$$

$$\frac{1}{\sin 1} \int \frac{\sin 1}{\cos(n-1) \cos(n-2)} du$$

$$= \frac{1}{\sin 1} \int \left[ \frac{\sin[(n-1)-(n-2)]}{\cos(n-1) \cos(n-2)} \right]$$

$$= \frac{1}{\sin 1} \int \frac{\sin(n-1) \cos(n-2) - (\cos(n-1) \sin(n-2))}{\cos(n-1) \cos(n-2)} du$$

$$= \frac{1}{\sin 1} \left[ \int \tan(n-1) du - \int \tan(n-2) du \right]$$

$$Q. 1) \int \frac{\sec^4 x}{\tan x} dx$$

$$= \int \frac{1 + \tan^2 x}{\tan x} \cdot 2k dk = \int \frac{1+k^2}{k} \cdot 2k dk =$$

$$= 2 \int (1+k^2) dk$$

$$\int (2 + 2t^4) dt = 2t + \frac{2t^5}{5} + C$$

$$\tan x = k^2$$

$$\sec^2 x dx = 2k dk$$

$$dx = \frac{2t dt}{\sec^2 x}$$

$$= 2 \int \left( \tan x + \frac{2}{5} \tan^5 x \right) dx = \frac{2t dt}{1 + \tan^2 x}$$

$$dx = \frac{2 \tan t}{1 - \tan^2 t}$$

$$dx = \frac{2t dt}{1 - t^2}$$

$$\rightarrow \frac{1}{\sec n} \cdot \sec n \cdot \tan n \cdot du = dp$$

$$(1) \int \frac{\ln^2(\sec n)}{\sec n} dn = \int p^2 dp$$

$$\int \frac{\ln(\sec n) \cdot \ln(\sec n) \cdot du}{\sec n}$$

$$\ln(\sec n) = p$$

$$\int \frac{dt}{t} \cdot \sqrt{t}$$

$$\text{Que } \int \frac{n \cos n}{(n \sin n + \cos n)^2} dn$$

$$n \sin n + \cos n = p$$

$$= \int \frac{dp}{p^2}$$

$$= n \cos n + \sin n - \sin n \cdot du = dp$$

$$\int \frac{dt}{t^2} = \int t^{-2} \cdot dt$$

$$= \frac{t^{-1}}{-1} + c = -\frac{1}{t} + c$$

⇒

$$(2) \int \frac{\sin 2n}{\sin n \cdot \sin 3n} dn = \frac{\sin(5n - 3n)}{\sin n \sin 3n}$$

$$= \frac{\sin n \cos 2n - \cos n \sin 2n}{\sin n \sin 3n}$$

$$= \int \frac{\cot 3n}{\sin n \sin 3n} dn - \int \frac{\cot n}{\sin n \sin 3n} dn$$

$$= \int \cot 3n \cdot dn - \int \cot n \cdot dn$$

$$= \frac{\ln(\sin 3n)}{3} - \frac{\ln(\sin n)}{1}$$

Ques:  $\int \frac{\sin 2n}{(a \sin^2 n + b \cos^2 n)^2} dn$       $a \sin^2 n + b \cos^2 n = k$   
 $= \frac{1}{(a-b)} \int \frac{dk}{k^2}$       $= (a-b) \int \sin 2n dn = dk$

(2)  $\int \frac{\ln^2 \left( \frac{n}{n+1} \right)}{n(n+1)} dn$       $= \log^2 n = (\log n)^2$

$\int t^2 dt$

$= \frac{1}{\left(\frac{n}{n+1}\right)} \cdot \frac{dn}{dn} \left(\frac{n}{n+1}\right) = dn \cdot dt$

$\Rightarrow \int t^2 dt$

$\int \frac{t^3}{3} + 1$

$\Rightarrow \frac{n+1}{n} \cdot \frac{(n+1)-n}{(n+1)^2} dn = dt$

$= \frac{\ln^3 \left( \frac{n}{n+1} \right)}{3} + 1$

$\frac{1}{n(n+1)} dn = dt$

\*  $\Rightarrow \int \frac{dn}{a \sin n + b \cos n}$


$\Rightarrow \frac{1}{\sqrt{a^2+b^2}}$   
 $a \sin n + b \cos n = \sqrt{a^2+b^2} \left( \frac{a}{\sqrt{a^2+b^2}} \sin n + \frac{b}{\sqrt{a^2+b^2}} \cos n \right)$

$= \frac{1}{\sqrt{a^2+b^2}} \int \frac{dn}{\cos(n-\phi)}$

$\left. \begin{aligned} &+ \frac{b}{\sqrt{a^2+b^2}} \cdot \cos n \\ &+ \frac{a}{\sqrt{a^2+b^2}} \cdot \sin n \end{aligned} \right\}$

$= \sqrt{a^2+b^2} [\sin n \cdot \cos \phi + \cos n \cdot \sin \phi]$

$= \sqrt{a^2+b^2} [\cos(n-\phi)]$



Ques  
(1)

$$\int \frac{du}{\sqrt{3} \cos u - \sin u}$$

$$\sqrt{3} \cos u - \sin u = \sqrt{(\sqrt{3})^2 + (-1)^2}$$

$$= 2 \cdot \left( \frac{\sqrt{3}}{2} \cos u - \frac{1}{2} \sin u \right)$$

$$\left( 2 \sin \frac{\pi}{3} \cos u - \sin u \left( \frac{\pi}{3} \right) \right)$$

$$2 \left( \cos u \left( \frac{\pi}{3} \right) - \sin u \sin \frac{\pi}{3} \right)$$

(2)

$$\int \frac{du}{\sin u + \cos u}$$

$$\frac{1}{\sqrt{2}} \cdot \int \frac{du}{\sin \left( u + \frac{\pi}{4} \right)}$$

$$\frac{1}{\sqrt{2}} \int \frac{du}{\cos \left( u - \frac{\pi}{4} \right)}$$

Ques:

$$\int \frac{2 du}{\sin u + \cos u} \quad \int \frac{2 du}{\sec u + \operatorname{cosec} u}$$

Ans:

$$\frac{2 du}{\frac{1}{\cos u} + \frac{1}{\sin u}} =$$

$$\frac{2 \sin u \cos u}{\sin u + \cos u} du$$

$$\int \frac{(1 + 2 \sin u \cos u) - 1}{\sin u + \cos u} = \int \frac{(\sin^2 u + \cos^2 u + 2 \sin u \cos u)}{\sin u + \cos u}$$

$$= \int \frac{(\sin u + \cos u)^2}{\sin u + \cos u} = \int (\sin u + \cos u) du = \int$$

$$\frac{1}{\sin u \cos u} du$$

$O-1 \Rightarrow 4, 6, 7, 8, 13$   
 Do first  $\rightarrow S-1 \Rightarrow 1, 2, 3, 9,$

Que:  $\int (\sin n - \cos n) (\sin n + \cos n)^5 dn$   
 $\int -k^5 dk$

$\sin n + \cos n = k$   
 $(\cos n - \sin n) dn = dk$

(2)  $\int \frac{\tan n + \sec n - 1}{\tan n - \sec n + 1} dn$

$\int \frac{\tan n + (\sec n - 1)}{\tan n + (\sec n - 1)} \times \frac{\tan n - (\sec n - 1)}{\tan n - (\sec n - 1)}$

$\int \frac{\tan n + \sec n - 1}{\tan n - \sec n + (\sec^2 n - \tan^2 n)}$

$\int \frac{\sec n + \tan n - 1}{(\sec n - \tan n) (\sec n + \tan n - 1)}$   
 $= \int (\sec n + \tan n) dn$

(3)  $\int \frac{\cos 2n}{\sin n} dn$

$\int \frac{1 - 2\sin^2 n}{\sin n} = \int \operatorname{cosec} n - 2 \int \sin n dn$

Ques  $\int \frac{dx}{e^x+1}$

Ans:  $\int \frac{dx}{e^x+1} = \int \frac{dt}{t+e^t}$

$$e^x = t$$

$$e^x dx = dt$$

$$dx = \frac{dt}{e^x}$$

$$= \int \frac{e^{-t}}{1+e^{-t}} = \int \frac{-dk}{k} \quad e^{-t} dt = -dk$$

$$= -\ln k$$

Ques:  $\int \frac{e^x-1}{e^x+1} dx$

$$e^x = t$$

$$e^x dx = dt$$

$$dx = \frac{dt}{e^x}$$

$$= \int \frac{e^x+1-2}{e^x+1} dx$$

$$= \int dx - \int \frac{2}{e^x+1} dx$$

$$\int dx - \int \frac{2e^{-x}}{1+e^{-x}} dx \quad [1+e^{-x}=k]$$

\* Integration of the form

\*  $\int \sin^m x \cos^n x dx$

1. if  $m = \text{odd}$  then put  $\cos x = t$

2. if  $n = \text{odd}$  put  $\sin x = t$

3. if  $n, m$  both are odd then put either of them  $t$ .

$$\text{QST} = t =$$

\* If both are even then use trigonometric identity to solve

\* otherwise, Put  $\tan x = t$  or  $\cot x = t$

Ex:  
Ex)  $\int \sin^3 x \cos^2 x \, dx$        $\cos x = t$

$$= \int \sin^2 x \cdot t^2 (-dt)$$

$$= \int (1 - \cos^2 x) (-t^2) \, dx$$

$$= \int (1 - t^2) (-t^2) \, dx$$

$$= \int (t^4 - t^2) \, dt$$

Q.  $\int \sin^5 x \cos x \, dx$

$$\int k^5 \, dk$$

$$\sin^5 x = t$$

$$\cos^5 x \, dx = t$$

$$dx = \frac{t}{\cos^5 x}$$

Q.  $\int \sin^4 x \, dx$

$$\sin x = k$$

$$\cos x \, dx = dk$$

Q.  $\int \sin^4 x \cos^2 x \, dx$

Q.  $\int \cos^4 x \, dx$

use trigonometric Id.

$$1) \int \sin^3 n \cos^{15} n \, dn \quad \text{---} \int$$

$$\int \frac{\sin^3 n}{(1-k^3)} \cdot k^{15} \cdot \frac{dn}{-2 \sin 2n} dk$$

$$\begin{aligned} \cos 2n &= k \\ -\sin 2n \, dn &= dk \\ dn &= \frac{k}{-\sin 2n} \end{aligned}$$

$$(2) \int \cos^3 n \sqrt{\sin n} \, dn$$

$$\int \frac{dt^3 \cdot dt}{\cos n}$$

$$\begin{aligned} \sin n &= t \\ \cos n \, dn &= dt \\ dn &= \frac{dt}{\cos n} \end{aligned}$$

$$(3) \int \cos^3 n \, dn$$

$$\textcircled{1} \int (1 - \cos^2 n) k^{15} \, dk$$

$$\begin{aligned} \cos n &= k \\ \sin n \, dn &= -dk \end{aligned}$$

$$\int (k^2 - 1) k^{15} \, dk$$

$$(2) \int (1 - \sin^2 n) \, dn = dt = \int (1 - t^2) \sqrt{t} \, dt \quad \left( \begin{array}{l} \sin n = t \\ \cos n \, dn = dt \end{array} \right)$$

$$\textcircled{3} \int (1 - \sin^2 n) \, dy = \int (1 - u^2) \, du$$

$$\begin{aligned} \sin n &= u \\ \cos n \, dn &= du \end{aligned}$$



Ques:  $\int (\sin n)^{-11/3} \cdot (\cos n)^{-1/3} dn$

$\frac{-11-1}{3} = \frac{-12}{3} = -4$

$= \frac{(\sin n)^{-11/3}}{\cos n^{1/3}} \frac{dn}{dn} = \int \frac{\tan n^{-4} dn}{\sec^2 n}$

Sin n = t

$(\tan n = t)$

$\sec^2 n dn = t$

$dn = \frac{t}{\sec^2 n}$

$\int \frac{1}{\sin n^{11/3}} \cdot \frac{1}{\cos n^{1/3}} dn$

$= \int \frac{t^{-3}}{\sec^2 n}$

Ans:  $\int \frac{(\sin n)^{-11/3} \cdot (\cos n)^{-1/3}}{(\cos n)^{-11/3}} \cdot (\cos n)^{-1/3} dn$

$\int (\tan n)^{-11/3} \cdot (\cos n)^{-4} dn =$

$\tan n = t$   
 $\sec^2 n dn = dt$

$= \int (\tan n)^{-11/3} \cdot \frac{1}{\cos^4 n} dn = \int (\tan n)^{-11/3} \cdot \sec^2 n \cdot \sec^2 n dn$

$= \int t^{-11/3} (1+t^2) dt$

Expression

$a^2 - x^2$

$a^2 + x^2$

$x^2 - a^2$

$\int \frac{a-x}{a+x} \text{ or } \int \frac{a+x}{a-x}$

$\int \frac{x}{a+x} \text{ or } \int \frac{a+x}{x}$

Sub

Put  $x = a \sin \theta$  or  $a \cos \theta$

$x = a \tan \theta$ ,  $x = a \cot \theta$

$x = a \sec \theta$ ,  $x = a \csc \theta$

$x = a \cos \theta$  or  $x = a \sin \theta$

$x^2 = a \tan^2 \theta$  or  $x = a \cot^2 \theta$

$$\int \frac{a-x}{x-b} \approx \int \frac{x-a}{a-x} \text{ or } \int \frac{(a-x)(\cos \theta - b)}{a-x} \quad \left| \quad x = a \cos^2 \theta + b \sin^2 \theta \right.$$

Ques.  $\int \sqrt{\frac{1+x}{1-x}} dx$

$$\int \frac{\cos \theta/2 \cdot (-2 \sin \theta/2 \cos \theta/2)}{\sin \theta/2} d\theta$$

$$x = \cos \theta \\ du = -\sin \theta d\theta$$

$$\frac{1+x}{1-x} = \frac{1+\cos \theta}{1-\cos \theta} = \frac{1+2\cos^2 \theta/2 - 1}{1-1+2\sin^2 \theta/2} = \cot^2 \theta/2$$

$$\int -2 \cos^2 \theta/2 d\theta = \int -( \cos \theta + 1 ) d\theta$$

Ques:  $\int \frac{1-\cos x}{\sin x - 2} dx$

$$1 \cos^2 x + 2 \sin^2 x = k$$

$$\text{or } 2 \cos^2 x + \sin^2 x = p$$

Ans:

$$(-\sin 2x + 2 \sin 2x)$$

Ques:  $\int \frac{1-x}{x-2} dx$

$$x \quad 2 \cos^2 x + 1 \sin^2 x = t$$

$$2(-\sin^2 x)$$

$$\cos^2 x \cdot 2(-\sin^2 x) + \sin^2 x \cdot \cos 2x du = dt$$

Answer

on Next Page.

$$\text{Ans} \int \sqrt{\frac{1-x}{x-2}} dx$$

$$x = 1 \cos^2 \theta + 2 \sin^2 \theta$$

$$dx = (-\sin 2\theta + 2 \sin 2\theta) d\theta$$

$$dx = \sin 2\theta d\theta$$

$$1-x = 1 - (\cos^2 \theta + 2 \sin^2 \theta)$$

$$= 1 - \cos^2 \theta - 2 \sin^2 \theta$$

$$= \sin^2 \theta - 2 \sin^2 \theta$$

$$= -\sin^2 \theta$$

$$x-2 = \cos^2 \theta + 2 \sin^2 \theta - 2$$

$$= \cos^2 \theta - 2(1 - \sin^2 \theta) = \cos^2 \theta - 2 + 2 \sin^2 \theta$$

$$= -\cos^2 \theta$$

$$= \int \sqrt{\frac{-\sin^2 \theta}{-\cos^2 \theta}} \times \sin 2\theta d\theta$$

$$= \int \frac{\sin \theta}{\cos \theta} \times 2 \sin \theta \cos \theta d\theta = \int (1 - \cos 2\theta) d\theta$$

$$\text{Ques:} \int \sqrt{\frac{x}{4+x}} dx$$

Ans

$$x = 4 \tan^2 \theta$$

$$4 \tan^2 \theta =$$

$$dx = 4 \sec^2 \theta + \tan^2 \theta$$

$$dx = 4 \cdot 2 \tan \theta \cdot \sec^2 \theta d\theta$$

$$\frac{x}{4+x}$$

$$= \frac{4 \tan^2 \theta}{4(1 + \tan^2 \theta)}$$

$$= \frac{\tan^2 \theta}{\sec^2 \theta}$$

$$= \sin^2 \theta$$

$$\int \sin \theta \times 8 \tan \theta \cdot \sec^2 \theta d\theta$$

$$\sin \theta \int \frac{\sin^2 \theta}{\cos^3 \theta} d\theta$$

Q.  $\int \sqrt{\frac{n}{1-n}} du = \int \sqrt{\frac{n}{1-n}} du.$   $n = \tan^2 \theta$

$$\int \sqrt{\frac{n}{1-n}} dn = \frac{\tan^2 \theta}{1 - \tan^2 \theta} = \frac{\tan^2 \theta}{\sec^2 \theta}$$

Ans:  $\int \frac{\sin^2 \theta}{\cos^2 \theta} \cdot 2 \sin \theta \cos \theta d\theta$   $n = \sin^2 \theta$   
 $dn = 2 \sin \theta d\theta$

$$\int \frac{\sin \theta}{\cos \theta} \cdot 2 \sin \theta \cos \theta d\theta$$

$$\int 2 \sin^2 \theta d\theta = \int (1 - \cos 2\theta) d\theta$$

Que:  $\int \cos(x) e^{2x} \sqrt{\frac{1-n}{1+n}} dn$

$n = \cos x$   
 $\cos x = n$   
 $\sin x$

$$\int 2e^{2x} \sqrt{\frac{1-\cos x}{1+\cos x}} \frac{1-\sin x}{1+\sin x} \times -dn$$

$\sin x = n$   
 $-\cos^2 x dx = dn$   
 $\cos^2 x dx = -dn$

$\int$

Ans:

$$n = \cos 2\theta$$
$$dn = -2 \sin 2\theta d\theta$$

$$\sqrt{\frac{1-n}{1+n}} = \sqrt{\frac{1-\cos 2\theta}{1+\cos 2\theta}} = \sqrt{\frac{2\sin^2 \theta}{2\cos^2 \theta}} = \tan \theta$$

$$\int \cos \left( 2 \cot^{-1} \sqrt{\frac{1-n}{1+n}} \right) dn$$

$$= \int \left( \cos 2 \cdot \cot^{-1} \left( \cot \left( \frac{\pi}{2} - \theta \right) \right) \right) dn$$

$$= \int \cos 2 \left( \frac{\pi}{2} - \theta \right) dn$$

$$= \int \cos (\pi - 2\theta) dn$$

$$= \int -\cos 2n \cdot (-2 \sin 2\theta d\theta) = \int \sin 2\theta d\theta$$

Ques:  $\int \sqrt{\frac{1-\sqrt{n}}{1+\sqrt{n}}} dn$

Ans:  $\sqrt{n} = \cos \theta$   
 $n = \cos^2 \theta$   
 $dn = -2 \sin \theta d\theta$

$$\frac{1-\sqrt{n}}{1+\sqrt{n}} = \frac{1-\cos \theta}{1+\cos \theta} = \frac{\sin^2 \theta/2}{\cos^2 \theta/2}$$

$$I = \int \sqrt{\frac{1-\sqrt{n}}{1+\sqrt{n}}} dn$$

$$= \int \frac{\sin \theta/2 - 2 \sin \theta \cos \theta}{\cos \theta/2} d\theta$$

$$= \int \frac{\sin \theta/2 - 2 \cdot 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} \cos \theta}{\cos \theta/2} d\theta$$

$$\cos \theta = 1 - 2 \sin^2 \frac{\theta}{2}$$

$$2 \sin^2 \frac{\theta}{2} = 1 - \cos \theta$$

$$= -2 \int (2 \sin^2 \frac{\theta}{2}) \cos \theta d\theta$$

Que:  $\int \frac{\sqrt{n}}{\sqrt{a^3 - n^3}} dn$

$$n^{3/2} = a^{3/2} \sin \theta$$

$$\frac{3}{2} n^{1/2} dn = a^{3/2} \cos \theta d\theta$$

$$= \frac{2}{3} \cdot a^{3/2} \int \frac{\cos \theta d\theta}{\cos \theta}$$

$$\boxed{n^{3/2} = a^{3/2} \cos \theta}$$

Que:  $\int \frac{n^2}{64 - n^6} dn = \int \frac{n^2}{\sqrt{2^6 - n^6}}$

$$n^3 = 2^3 \sin \theta$$

$$3n^2 dn = 8 \cos \theta d\theta$$

$$= \int \frac{8 \cos \theta d\theta}{2^3 \cdot \cos \theta}$$

Que:  $\int \frac{\sin n \, dn}{\sqrt{9 - \sin^4 n}}$

$\sin^2 n = k$   
 $\sin 2n \, dn = dk$

$$\int \frac{dk}{\sqrt{9 - k^2}}$$

$$= \sin^{-1} \left( \frac{k}{3} \right) + C$$

Que:  $\int \frac{e^n \, dn}{\sqrt{e^{2n} - 1}}$

$e^n = k$   
 $e^n \, dn = dk$

$$\int \frac{dk}{\sqrt{k^2 - 1}} = \ln |k + \sqrt{k^2 - 1}| + C$$

Que:  $\int \frac{e^n}{4 + e^{2n}} \, dn$

$e^n = t$   
 $e^n \, dn = dt$

$$\int \frac{dt}{2^2 + t^2} = \frac{1}{2} \tan^{-1} \frac{e^n}{2} + C$$

Que:  $\int \frac{dn}{\sqrt{2n - n^2 - 1}} = \int \frac{dn}{\sqrt{1 - (n-1)^2}} = \sin^{-1} (n-1)$

1. take common n coefficient
2. Rearrange +1 on +ve

## Integral of Rational functions (without partial fraction)

(1)  $\int \frac{1}{a} dx, \int a dx, \int \frac{1}{\sqrt{a}}$  -  $a = ax^2 + bx + c$

$$d. \int \frac{dx}{2x^2 - 3x + 4} = \frac{2x^2 - 3x + 4}{2} \cdot \left[ x^2 - \frac{3}{2}x + 2 \right]$$

$$= \frac{1}{2} \int \frac{dx}{\left(x - \frac{3}{4}\right)^2 + \left(\frac{\sqrt{23}}{4}\right)^2} = \frac{1}{2} \left[ x^2 - 2\left(\frac{3}{4}\right)x + \left(\frac{3}{4}\right)^2 + 2 - \left(\frac{3}{4}\right)^2 \right]$$

$$= \frac{1}{2} \left[ \left(x - \frac{3}{4}\right)^2 + \frac{23}{16} \right]$$

$$= \frac{1}{2} \cdot \frac{1}{\frac{\sqrt{23}}{4}} \cdot \tan^{-1} \left( \frac{x - \frac{3}{4}}{\frac{\sqrt{23}}{4}} \right) + C$$

$$= \frac{1}{2} \cdot \frac{1}{\frac{\sqrt{23}}{4}} \cdot \tan^{-1} \left( \frac{x - \frac{3}{4}}{\frac{\sqrt{23}}{4}} \right) + C$$

d.  $\int \frac{dx}{\sqrt{2x^2 - 3x + 4}} = \frac{1}{2} \int \frac{dx}{\left(x - \frac{3}{4}\right)^2 + \left(\frac{\sqrt{23}}{4}\right)^2}$

Formula No-18



$$\frac{A}{B} = 1 \cdot A = 1$$

Ques:  $\int \sqrt{3n^2 - 6n + 10} \, dn$

$$= \int \sqrt{3n^2 - 6n + 10} \, dn$$

$$3n^2 - 6n + 10$$

$$3 \left[ n^2 - 2n + 1 - 1 + \frac{10}{3} \right]$$

$$= \sqrt{3} \int \sqrt{(n-1)^2 + \left(\frac{\sqrt{7}}{3}\right)^2} \, dn$$

$$3 \cdot \left[ n^2 - 2n + \frac{10}{3} \right]$$

$$3 \cdot \left[ n^2 - 2n + 1 - 1 + \frac{10}{3} \right]$$

$$= \sqrt{3} \left( \frac{n-1}{2} \sqrt{(n-1)^2 + \frac{7}{3}} \right)$$

$$= 3 \left[ (n-1)^2 + \frac{7}{3} \right]$$

Type

(Q)  $\int \frac{L \, dn}{Q}, \int \frac{L}{\sqrt{Q}} \, dn, \int L \sqrt{Q} \, dn.$   $Q = an^2 + bn + c$   
 $L = pn + q$

Ques:  $L = \lambda \frac{dQ}{dn} + u$  Here  $\lambda = \text{constant}$

$$pn + q = \lambda(2an + b) + u$$

$$p = 2a\lambda -$$

$$q = \lambda b + u -$$

then move to type 1.

$$\text{Sol: } \int \frac{5n+7}{2n^2-3n+4} dn$$

$$5n+7 = \lambda \frac{d}{dn} (2n^2-3n+4) dn$$

$$5n+7 = \lambda (4n-3) + \mu$$

$$5n = 4n\lambda$$

$$\lambda = \frac{5n}{4n} = \frac{5}{4}$$

$$7 = -3\lambda + \mu$$

$$\mu = 7 + 3 \cdot \frac{5}{4} = \frac{43}{4}$$

$$= \int \frac{\frac{5}{4}(4n-3) + \frac{43}{4}}{2n^2-3n+4} dn$$

$$= \frac{5}{4} \int \frac{4n-3}{2n^2-3n+4} dn + \frac{43}{4} \int \frac{dn}{2n^2-3n+4}$$

↓

$$2n^2-3n+4 = p$$

$$(4n-3)dn = dp$$

$$= \frac{5}{4} \int \frac{dp}{p}$$

$$= \frac{5}{4} \ln(2n^2-3n+4) + \dots$$

Ques:  $\int \frac{3 \cos n + 5}{\sqrt{\sin^2 n - 2 \sin n + 5}} dn$

Ques:  $\int \frac{\frac{3}{2} \sin 2n + 5 \cos n}{\sqrt{\sin^2 n - 2 \sin n + 5}} dn$

$\sin n = t$   
 $\cos n dn = dt$

$= \int \frac{(3 \sin n + 5) \cos n}{\sqrt{\sin^2 n - 2 \sin n + 5}} dn$

$\int \frac{3t + 5}{\sqrt{t^2 - 2t + 5}} dt = \frac{3}{2}$

$\frac{3}{2} \int \frac{2t - 2}{\sqrt{t^2 - 2t + 5}} dt + 8 \int \frac{dt}{\sqrt{t^2 - 2t + 5}}$

$\frac{3}{2} \int \frac{2k dk}{k} + 8 \int \frac{dt}{2^2 + (t-1)^2} \quad \left| \quad \begin{aligned} 3t + 5 &= \frac{d}{dt}(t^2 - 2t + 5) + \mu \\ &= \frac{d}{dt}(2t - 2) + \mu \end{aligned} \right.$

$= \frac{3k + \mu}{\sqrt{t^2 - 2t + 5}}$

$3 = 2\mu \Rightarrow \mu = \frac{3}{2}$

$\mu = 5 + 2A$

$= 5 + 2 \times \frac{3}{2} = 8$

$= \int \frac{2t - 2}{\sqrt{t^2 - 2t + 5}} dt + 8 \int \frac{dt}{\sqrt{(t-1)^2 + 4}}$

$= \frac{3}{2} \int \frac{2k dk}{k} + 8 \int \frac{dt}{\sqrt{2^2 + (t-1)^2}}$

$= 3k + 8$

Type = 3

$$\int \frac{Q_1}{Q_2} dx, \int \frac{Q_1}{Q_2} dx$$

$$Q_1 = an^2 + bn + c$$

$$Q_2 = pn^2 + qn + r$$

$$Q_1 = d \cdot Q_2 + u \frac{d}{dx}(Q_2) + v.$$

{ Attemptly Question goes to  
type 1 and type 2 }

get the value of  $d, u$  and  $v$

$d, u, v$  are constant and balances.

Ques!  $\int \frac{3x^2 - 2x + 5}{x^2 - 2x + 10} dx$   $\frac{a}{a}$

$$3x^2 - 2x + 5 = d(x^2 - 2x + 10) + u(2x - 2) + v.$$

$$x = 3, \quad -2 = -2d + 2u$$

$$u = 2, \quad v = -21$$

$$\int \frac{3x^2 - 2x + 5}{x^2 - 2x + 10} dx$$

$$3 \int \frac{x^2 - 2x + 10}{x^2 - 2x + 10} dx + 2 \int \frac{2x - 2}{x^2 - 2x + 10} dx - 21 \int \frac{dx}{(x^2 - 2x + 10) + 9}$$

$$= 3n + 2 \ln(n^2 - 2n + 10) - 21 \cdot \frac{1}{3} \cdot \tan^{-1}\left(\frac{n-1}{3}\right)$$

Ans

## \* Integration of Trigonometric function!

! Type 1:

$$\int \frac{dn}{a + b \sin^2 n}, \quad \int \frac{dn}{a + b \cos^2 n}, \quad \int \frac{dn}{a \sin^2 n + b \cos^2 n + c}$$

$$\int \frac{dn}{a \sin^2 n + b \cos^2 n + c \sin n \cos n}$$

Trick! Const. is replaced by

$$(1) \quad a \rightarrow a(\sin^2 n + \cos^2 n)$$

(2) take  $\cos^2 n$  common from denominator and put  $\tan n = t$ .

Que:  $\int \frac{dn}{3 + 4 \sin^2 n} = \int \frac{dn}{3(\sin^2 n + \cos^2 n) + 4 \sin^2 n} =$

$$\int \frac{\frac{dn}{\cos^2 n}}{7 \sin^2 n + 3 \cos^2 n} = \int \frac{\sec^2 n \cdot dn}{7 \tan^2 n + 3}$$

$$= \frac{1}{7} \int \frac{\sec^2 n \cdot dn}{\tan^2 n + \left(\frac{\sqrt{3}}{7}\right)^2} = \frac{1}{7} \int \frac{dk}{k^2 + \left(\frac{\sqrt{3}}{7}\right)^2}$$

$$\text{Ques: } \int \frac{dx}{3 + \cos^2 x} = \int \frac{dx}{3 \sin^2 x + 4 \cos^2 x}$$

$$= \int \frac{\sec^2 x dx}{3 \tan^2 x + 4} = \int \frac{dt}{3t^2 + 4} \quad \tan x = t$$

$$\text{Ques: } \int \frac{dx}{3 \sin^2 x + 4 \cos^2 x - 8 \sin x \cos x}$$

$$= \frac{dx}{3 \sin^2 x} \int \frac{dt}{3t^2 - t + 4}$$

Type: 2  $\int \frac{dx}{a + b \sin^2 x}$ ,  $\int \frac{dx}{a + b \cos^2 x}$ ,  $\int \frac{dx}{a \sin x + b \cos x + c}$

$$a \rightarrow a \left( \cos^2 \frac{x}{2} + \sin^2 \frac{x}{2} \right)$$

$$\cos x \rightarrow \cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}$$

$$\sin x \rightarrow 2 \sin \frac{x}{2} \cos \frac{x}{2}$$

take  $\cos^2 \frac{x}{2}$  common and put  $\tan \frac{x}{2} = t$

$$J.A \Rightarrow 2,5$$

Ques!  $\int \frac{dn}{5+4 \sin n}$

$$\frac{5+4 \sin n}{5 \sin^2 \frac{n}{2} + 5 \cos^2 \frac{n}{2} + 8 \sin \frac{n}{2} \cos \frac{n}{2}}$$

Ans  $\int \frac{\sec^2 \frac{n}{2} dn}{5 + \tan^2 \frac{n}{2} + 8 \tan \frac{n}{2} + 5}$

$$\tan \frac{n}{2} = p$$

$$\sin^2 \frac{n}{2} dn = 2 dp$$

$$= \int \frac{2 dp}{5p^2 + 8p + 5}$$

Ques!  $\int \frac{dn}{3+2 \sin n + \cos n}$

$$\frac{3+2 \sin n + \cos n}{3 \sin^2 \frac{n}{2} + \dots}$$

$$= \int \frac{dn}{3(\cos^2 \frac{n}{2} + \sin^2 \frac{n}{2}) + \dots}$$

Ans!  $\int \frac{3(\sin^2 \frac{n}{2} + \cos^2 \frac{n}{2}) + 2 \cdot 2 \sin \frac{n}{2} \cos \frac{n}{2} + (\cos^2 \frac{n}{2} - \sin^2 \frac{n}{2})}{4 \cos^2 n + 2 \sin^2 n + 4 \sin n \cos n}$

$$= \frac{1}{2} \int \frac{\sec^2 \frac{n}{2} dn}{\tan^2 \frac{n}{2} + 2 \tan \frac{n}{2} + 2}$$

$$\tan \frac{n}{2} = k \quad = \int \frac{2 \cdot dk}{k^2 + 2k + 2}$$

24/06/17

Ques 1  $\int \frac{dn}{3\cos^2 n + 4\sin^2 n - 2}$        $dx$

$3\cos^2 n + 4\sin^2 n - 2\sin^2 n - 2\cos^2 n$

$\tan n = t$

$$\int \frac{dn}{\cos^2 n + 2\sin^2 n} = \int \frac{\sec^2 n dn}{2\tan^2 n + 1} = \int \frac{dt}{2t^2 + 1}$$

Q.  $\int \frac{dn}{1 + 2\sin n + 4\cos n}$

$$= \int \frac{\sin^2 \frac{n}{2} + \cos^2 \frac{n}{2} + 4\sin \frac{n}{2} \cos \frac{n}{2} + 4\cos^2 \frac{n}{2} - 4\sin^2 \frac{n}{2}}{2}$$

$$= \int \frac{5\cos^2 \frac{n}{2} + 4\sin \frac{n}{2} \cos \frac{n}{2} - 3\sin^2 \frac{n}{2}}{2}$$



Type 3

$$\int \frac{a \sin n + b \cos n + c}{p \sin n + q \cos n + r} dx.$$

Tech:

$$a \sin n + b \cos n + c = \lambda (p \sin n + q \cos n + r) + \mu \frac{d}{dn} (p \sin n + q \cos n + r) + v.$$

Compare coefficient of  $\sin n, \cos n$

and get the value of  $\lambda, \mu$  and  $v$ .

Ques:  $\int \frac{3 \sin n + 2 \cos n + 1}{\cos n + 2 \sin n - 3} dx$

Ans:

$$3 \sin n + 2 \cos n + 1 = \lambda (\cos n + 2 \sin n - 3) + \mu (\sin n + 2 \cos n) + v$$

$$= (3 \sin n + 2 \cos n + 1) = \lambda (2 \sin n + \cos n - 3) + \mu (-\sin n + 2 \cos n) + v$$

$$3 = 2\lambda - \lambda \quad \text{--- (1)} \quad \lambda = 8/5$$

$$2 = \lambda + 2\mu \quad \text{--- (2)} \quad \mu = 1/5$$

$$1 = -3\lambda + v \quad \text{--- (3)} \quad v = 29/5$$

$$\int \frac{3 \sin n + 2 \cos n + 1}{\cos n + 2 \sin n - 3} dx =$$

$$\frac{8}{5} \int dx + \frac{1}{5} \int \frac{-\sin x + 2 \cos x}{\cos x + 2 \sin x - 3} dx + \frac{29}{5} \int \frac{dx}{\cos x + 2 \sin x - 3}$$

$$= \frac{8}{5} x + \frac{1}{5} \ln(\cos x + 2 \sin x - 3) + \dots$$

and geses type 2.

Qu:  $\int \frac{6 + 3 \sin x + 14 \cos x}{3 + 4 \sin x + 5 \cos x} dx$

$$6 + 3 \sin x + 14 \cos x = A(3 + 4 \sin x + 5 \cos x) + B(4 \cos x + 5 \sin x) + C$$

$$6 = A \cdot 3$$

$$3 = 3A =$$

$$A = \frac{6}{3} = 2$$

$$3 = 4B - 5C$$

$$-4 = 3, \quad C = -3$$

$$B = 1$$

$$14 = (4B - 5C) + C$$

$$14 = 4 + 4 = 14 - 3 = 11$$

$$C = 0$$

$$A = 2, \quad B = 1, \quad C = 0$$

$$= 2 \int \frac{3 + 4 \sin x + 5 \cos x}{3 + 4 \sin x + 5 \cos x} dx + \int \frac{4 \cos x - 5 \sin x}{3 + 4 \sin x + 5 \cos x} dx$$

$$= 2x + \ln(3 + 4 \sin x + 5 \cos x) + 1$$

Ques<sup>o</sup>  $\int \frac{3e^n + 5e^{-n}}{4e^n - 5e^{-n}} dn$

~~$= \int 3e^n$~~

$$3e^n + 5e^{-n} = \lambda(4e^n - 5e^{-n}) + \frac{B}{4e^n - 5e^{-n}} + \frac{C}{1}$$

~~Part =  $\frac{3e^n + 5e^{-n}}{4e^n - 5e^{-n}}$~~   
 ~~$A = \frac{3e^n + 5e^{-n}}{4e^n - 5e^{-n}}$~~

$3 = 4A + B$  — (i)

$5 = -5A + 5B$  — (ii)

$A = -\frac{1}{8}, B = \frac{7}{8}$

$= A \int dn + \frac{7}{8} \int \frac{4e^n + 5e^{-n}}{4e^n - 5e^{-n}} dn$

$= -\frac{1}{8}n + \frac{7}{8} \ln(4e^n - 5e^{-n}) + C$

Q.  $\Rightarrow \int \frac{\sin^n}{e^n - \sin^n - \cos^n} dn$

$= \int \frac{e^{-n} \sin^n dn}{1 - (\sin^n + \cos^n) e^{-n}}$

$= \frac{1}{2} \int \frac{dt}{1-t}$

$(\sin^n + \cos^n) e^{-n} = t$


$(\cos^n - \sin^n) e^{-n} - e^{-n} (\sin^n + \cos^n) \times dn = dt$

$-2 \sin^n e^{-n} dn = dt$

$\sin^n e^{-n} dn = -\frac{1}{2} dt$

# \* Integration using Partial fraction:

$$\int \frac{P(x)}{Q(x)} dx$$



Where P and Q both are polynomial & degree of P < degree of Q

- In partial fraction Q(x) can be
- (i) Linear and non repeated.
  - (ii) Linear and repeated
  - (iii) Quadratic and non-repeated
  - (iv) Quadratic and repeated.

Ex:  $\int \frac{dx}{(x-1)(2x-3)}$

$$\frac{1}{(x-1)(2x-3)} = \frac{A}{x-1} + \frac{B}{2x-3} \quad \text{when } A, B \text{ are constants}$$

Partial fraction  
3rd degree and 2nd degree  
& Degree of numerator < Denominator

$$1 = A(2x-3) + B(x-1) \quad (\text{LCM}) \quad (\text{An Identity})$$

$$0 = 2A + B \quad \text{--- (i)}$$

$$1 = -3A - B \quad \text{--- (ii)}$$

Let  $x = \frac{3}{2}$

$$1 = B \cdot \left(\frac{3}{2} - 1\right) = B = 2$$

$$A = -1$$

$$= -\int \frac{dy}{n-1} + 2 \int \frac{dy}{2n-3}$$

$$= -\ln(n-1) + 2 \frac{\ln(n-3)}{2} + C$$

Ques:

$$\int \frac{dx}{(x-1)^2(x+2)}$$

$$= \frac{px+q}{(x-1)^2} + \frac{r}{x+2}$$

$$\rightarrow \frac{A}{(x-1)} + \frac{B}{(x-1)^2} + \frac{C}{(x+2)} \quad \text{--- (let)}$$

$$= \frac{A(x-1) + B}{(x-1)^2} + \frac{C}{(x+2)}$$

$$= \frac{Ax + (B-A)}{(x-1)^2} + \frac{C}{(x+2)}$$

$$\frac{A}{(x-1)} \quad A(x-1) + B$$

$$\frac{A}{(x-1)} + \frac{B}{(x-1)^2} + \frac{C}{(x+2)}$$

$$1 = A(x-1) + B + C(x+2)$$

$$x=1 \quad B = \frac{1}{3}$$

$$\rightarrow x=-2 \quad C = \frac{1}{9}$$

$$x=0 \quad 1 = -2A + 2B + C$$

$$2A + 2A = 2B + C - 1$$

$$= \frac{2}{3} + \frac{1}{9} - 1$$

$$\frac{6 + 1 \cdot 9}{9} = \frac{2}{9} \quad \left\{ A = -\frac{1}{9} \right\}$$

$$\int \frac{dx}{x} = \frac{1}{9} \int \frac{dx}{x-1} + \frac{1}{3} \left( \frac{dx}{(x-1)^2} + \frac{1}{9} \left( \frac{dx}{x+2} \right) \right)$$

$$= -\frac{1}{9} \ln(x-1) + \frac{1}{3} \cdot \left( -\frac{1}{x-1} \right) + \frac{1}{9} \ln(x+2)$$

$$\underline{\underline{\text{Ans}}} = \frac{1}{(x-1)(x+2)(2x+3)}$$

$$= \frac{A}{x-1} + \frac{B}{x+2} + \frac{C}{2x+3}$$

$$1 = A(x+2)(2x+3) + B(x-1)(2x+3) + C(x+1)(x+2)$$

$$\begin{aligned} A \cdot 1 &= A + 2 + B + 2 + C \\ 1 &= 2A + 3A + -B + 3B + C + 2C \end{aligned}$$

$$\text{Ans!} \quad x = 1 \quad A = \frac{1}{15}$$

$$x = -2 \quad B = \frac{1}{3}$$

$$x = -\frac{3}{2} \quad C = \frac{4}{5}$$

$$= \frac{1}{15} \ln(n-1) + \frac{1}{3} \ln(n+2) - \frac{4}{5} \ln\left(\frac{2n+3}{2}\right) + C$$

$$\begin{aligned} Q. \int \frac{dx}{(x-1)(x+2)} &= \frac{1}{3} \left[ \frac{dx}{x-1} - \frac{dx}{x+2} \right] \\ &= \frac{1}{3} (\ln|x-1| - \ln|x+2|) \end{aligned}$$

$$Ques! \int \frac{x+5}{(x-2)^2} dx$$

$$= \frac{x+5}{(x-2)^2} = \frac{A}{x-2} + \frac{B}{(x-2)^2}$$

$$\begin{aligned} x+5 &= A(x-2) + B \\ x=2, \quad B &= 7 \\ x=0, \quad A &= 1 \end{aligned}$$

$$\int \frac{dx}{x-2} + 7 \int \frac{dx}{(x-2)^2}$$

$$= \ln|x-2| - \frac{7}{x-2} + C$$

$$Ques! \int \frac{dx}{(x^2+1)(x^2+2)}$$

$$\int \frac{dx}{x^2+1} = \int \frac{dx}{x^2+2}$$

$$= \tan^{-1} x - \frac{1}{\sqrt{2}} \tan^{-1} \frac{x}{\sqrt{2}} + C$$

Que:  $\int \frac{dx}{(\sin x - \sin^2 x)}$

$$\int \frac{dx}{\sin x (1 - 2 \cos x)} = \int \frac{\sin x dx}{\sin^2 x (1 - 2 \cos x)}$$

$$\int \frac{\sin x dx}{(1 - \cos^2 x)(1 - 2 \cos x)}$$

$$= \int \frac{\sin x dx}{(2 \cos x - 1)(\cos x - 1)(\cos x + 1)}$$

$$= - \int \frac{dt}{(2t-1)(t-1)(t+1)}$$

Que:  $\int \frac{x^2 + 7}{(x+1)(x^2+4)}$

Ans:  $\frac{A}{x+1} + \frac{Bx+C}{x^2+4}$

$$x^2 + 7 = A(x^2 + 4) + (Bx + C)(x + 1)$$

$$x = -1$$

$$A = 1$$

$$2 = A + B$$

$$B = -1$$

$$C = 3$$

$$5 = A(1+4) + (Bx+C)(-1+1)$$

$$A = \frac{5}{5} = 1$$



# \* Integration by Parts!


$$\int f(x) \cdot g(x) dx = f(x) \cdot \int g(x) dx - \int \left( \frac{d}{dx} f(x) \right) \cdot g(x) dx$$

1. 1st  $\&#947$   $f(x)$  is diff. easily.  
2nd easily integrable

Ex!  $\int \underset{\text{I}}{x} \sin x dx = x \int \underset{\text{II}}{\sin x} dx - \int \left( \frac{d}{dx} x \right) \cdot (\sin x) dx$

$$= -x \cos x - \int 1 \cdot (-\cos x) dx$$
$$= -x \cos x + \int \cos x dx$$
$$= -x \cos x + \sin x + C$$

Some times ILATE Rule is helpful to choose in decreasing order



$$\int_{I} f g' du = f g(u) - \int (f' g(u)) du$$

$$\int \overset{\text{I}}{u} \overset{\text{II}}{\sin u} du$$

$\downarrow$                        $\downarrow$   
 A                          T

$$\int \overset{\text{II}}{1} \cdot \overset{\text{I}}{\sin^{-1} u} du$$

$$\int \overset{\text{II}}{e^u} (u^2 - 2u + 4) \overset{\text{I}}{du}$$

$\downarrow$                        $\downarrow$   
 Exp                      Ag.

$$\int \overset{\text{I}}{\ln u} \overset{\text{II}}{\cos u} du$$

Que:  $\int \overset{\text{I}}{u} \overset{\text{II}}{\tan^{-1} u} \overset{\text{I}}{du}$

(2)  $\int \overset{\text{II}}{u} \overset{\text{I}}{e^u} du$

$$= \rightarrow u \int \tan^{-1} u - \int \left( \frac{d}{du} u \right) \tan^{-1} u du$$

$$= (\tan^{-1} u) \frac{u^2}{2} - \int \frac{u^2}{1+u^2} du$$

$$= \frac{u^2}{2} \tan^{-1} u - \frac{1}{2} \int \frac{(u^2+1) - 1}{1+u^2} du$$

$$= \frac{u^2}{2} \tan^{-1} u - \frac{1}{2} \left( 1 - \frac{1}{1+u^2} \right) du = \frac{u^2}{2} \tan^{-1} u - \frac{1}{2} [u - \tan^{-1} u]$$

(2)  $\int \overset{\text{I}}{u} \overset{\text{II}}{e^u} du$

$$= u \int e^u du - \int \left( \frac{d}{du} u \right) \cdot e^u du$$

$$= u e^u - \int 1 \cdot e^u du$$

$$= u e^u - e^u$$

$$\text{Qw1)} \int \frac{\operatorname{cosec}^2 n}{\text{II}} \cdot \ln(\sec n) \text{ I} \, dn$$

$$\operatorname{cosec}^2 n \int \frac{1}{\sec n} + \int (\operatorname{cosec}^2 n \sec n - \int \sec n \, dn)$$

$$\operatorname{cosec}^2 n$$

$$= \ln(\sec n) (-\cot n) - \int \frac{1}{\sec n} \sec n \operatorname{tann} \times (-\cot n) \, dn$$

$$= -\cot n \ln(\sec n) + n + C$$

$$\text{Qw' 1)} \int \frac{\sin n}{\text{II}} \ln(\sec n + \operatorname{tann}) \text{ I} \, dn$$

$$\ln(\sec n + \operatorname{tann}) = \operatorname{cosec} n + \int \operatorname{cosec} n \cdot \frac{1}{\sec n + \operatorname{tann}} \operatorname{cosec}^2 n \, dn$$

$$\ln(\sec n + \operatorname{tann}) = \operatorname{cosec} n - \sin n \cdot \frac{1}{\sec n + \operatorname{tann}} \int dn$$

Imp.

(2)  $\frac{\cos^{-1} x}{x^3} dx$

$\cos^{-1} x = \alpha$   
 $x = \cos \alpha$   
 $dx = -\sin \alpha d\alpha$

$\frac{\cos^{-1} \cos \alpha}{\cos^3 \alpha} =$

$\int \frac{\alpha - \sin \alpha d\alpha}{\cos^3 \alpha} = \int \alpha \left( \frac{\tan \alpha \cdot \sec^2 \alpha}{\cos^3 \alpha} \right) d\alpha$

$= -\alpha \cdot \frac{1}{2} \tan^2 \alpha + \frac{1}{2} \int 1 \cdot \tan^2 \alpha d\alpha$

$= -\frac{\alpha}{2} \tan^2 \alpha + \frac{1}{2} \int (\sec^2 \alpha - 1) d\alpha$

$= -\frac{\alpha}{2} \tan^2 \alpha + \frac{1}{2} [\tan \alpha - \alpha] + C$

Stav  $\sec^2 \alpha d\alpha = k$   
 $= \int k dk = \frac{k^2}{2}$

$\tan \alpha = k$

$\sec^2 \alpha d\alpha = dk$

Q.1  $\int x (\sin x \cos^2 x) dx$

$\Rightarrow -\frac{x \cos^3 x + 1}{3} \int \cos^3 x dx$

$\int \sin x \cos^2 x dx$   
 $= -\int t^2 dt$   
 $= -\frac{t^3}{3}$   
 $= -\frac{1}{3} \cos^3 x$

$\cos x = t$

$\sec^2 x dx = dt$

Q.2  $\int \ln(x + \sqrt{x^2 + 1}) dx$

Ans

$$x \ln(x + \sqrt{x^2+1}) - \int \frac{x}{\sqrt{x^2+1}} dx$$

$$= - \int \frac{p dp}{p - p + 1} \\ = - \int \frac{p dp}{1}$$

$$1 + \frac{1 \cdot 2x}{2\sqrt{x^2+1}} \\ = \frac{1 + \frac{2x}{2\sqrt{x^2+1}}}{1 + \frac{2x}{2\sqrt{x^2+1}}}$$

$$= \frac{\sqrt{x^2+1} + x}{1 + \frac{2x}{2\sqrt{x^2+1}}} \cdot \frac{1}{\sqrt{x^2+1}}$$

$$= \frac{1}{\sqrt{x^2+1}}$$

$$x^2+1 = p^2$$

$$x dx = p dp$$

Q.  $\int_I x^2 e^x dx - \int_{II} x e^x dx$

$$x^2 e^x dx - \int (x \cdot \int e^x dx) dx$$

$$x^2 e^x - 2 \int_I x e^x dx = x^2 e^x - 2(n-1)e^{n+1} \quad \text{Ans}$$

↓  
J

$$J = \int_I x e^x$$

$$= x e^x - \int 1 \cdot e^x dx$$

$$= x e^x - e^x$$

$$= (x-1)e^x$$

Ans.

\* Two classic Integrations!

$$\int e^n (f(n) + f'(n)) dn = e^n f(n) + C$$

$$\int (f(n) + n f'(n)) = n f(n) + C$$

$$Q. \int e^n f(n) dn + \int e^n f'(n) dn$$

$$= \int e^n f(n) dn + e^n f(n) - \int e^n f(n) dn$$

$$Q_{2} \int e^n \left( \underset{f}{\sin n} + \underset{f'}{\cos n} \right) dn = e^n \sin n + C$$

$$Q_{3} \int e^n \left[ \underset{f'}{\sec^2 n} + \underset{f}{\tan n} \right] dn = e^n \sec^2 n \\ = e^n \tan n$$

$$Q. \int e^n [\cot n + \operatorname{cosec}^2 n] dn = e^n \cot n$$

$$Q. \int \left( \underset{f}{2 \sin n} + n \underset{f'}{\cos n} \right) dn = n \sin n + C$$

S.M ⇒ 5 class

How: Q.1, 3 both both side.

14, Q.9 classic Int.

8-1 ⇒ Q. 6, 7, 8, 10, 11, 17,

$$\text{Ques: } \int \frac{x e^x}{(1+x)^2} dx$$

$$= \int e^x \left[ \frac{x+1-1}{(x+1)^2} \right] dx$$

$$= \int e^x \cdot \left[ \frac{1}{1+x} - \frac{1}{(1+x)^2} \right] dx$$

$$= \int e^x \cdot \left[ \frac{1}{\underset{f}{1+x}} - \frac{1}{\underset{f'}{(1+x)^2}} \right] dx$$

$$= \frac{e^x}{1+x} + \cot x$$

\* Two Classic Integrals

$$\int e^n [f(n) + f'(n)] dn = \int e^n (f(n) + 1)$$

$$\int e^n f(n)$$

\*  $\int e^n [f(n) - f''(n)] dn = \int e^n [f(n) + f'(n) - f'(n) - f''(n)] dn$

$$\int e^n [f(n) + f'(n)] dn - \int e^n [f'(n) + f''(n)] dn$$

$$= \int e^n f(n) - \int e^n [f'(n) + f''(n)] dn$$

$$= \int e^n f(n) - \int e^n f'(n)$$

$$= \int e^n (f(n) - f'(n)) dn$$

Ex  $\int \frac{n + \sin n}{1 + \cos n} dn$

$$= \int \frac{n + 2 \sin \frac{n}{2} \cos \frac{n}{2}}{1 + 2 \cos^2 \frac{n}{2} - 1} dn = \int \left( \frac{n}{2} + \frac{n}{2} - \frac{1}{2} \sec^2 \frac{n}{2} \right) dn$$

$$= \frac{n + \frac{n}{2}}{2} + 1$$



$$(2) \int \frac{e^n (n-1)}{(n+1)^3}$$

$$\int \frac{e^n (n+1-2)}{(n+1)^3} = \int e^n \left[ \frac{1}{n(n+1)^2} - \frac{2}{(n+1)^3} \right] dx$$

$$= \int e^n \left[ \frac{1}{1+n^2} + \frac{-2}{1+n} \right] dx$$

$$= e^n \left[ \frac{1}{1+n} + \dots \right]$$

Ques:  $\int e^n [\ln(\sec n + \tan n) + \sec n] dx$   
 $= e^n \ln(\sec n + \tan n) + \dots$

$$(2) \int \frac{e^n (n^2+1)}{(n+1)^2} dx$$

$$= \frac{e^n (n^2-1+2)}{(n+1)^2} = e^n \left[ \frac{(n-1)(n+1) + 2}{(n+1)^2} \right]$$

$$\int e^n \left[ \frac{n+1}{n+1} + \frac{2}{(n+1)^2} \right] = \left[ \frac{n+1}{n+1} \right] e^n$$

$\uparrow$                        $\uparrow$   
 $f$                        $f'$

Ques.  $\int (\sin(\ln x) + \cos(\ln x)) dx$        $\ln x = t$   
 $x = e^t$   
 $dx = e^t dt$

$$I = \int (\sin t + \cos t) e^t dt$$

$$= e^t \sin t + t$$

M-2  $\int (\sin(\ln x) + \frac{1}{x} \cos(\ln x)) dx$

$\uparrow$                        $\uparrow$                        $\uparrow$   
 $f'$                        $\frac{1}{x}$                        $f$

M  $e^{\tan^{-1} x} \frac{(1+x+x^2)}{1+x^2}$

$\tan^{-1} x = \theta$   
 $x = \tan \theta$   
 $dx = \sec^2 \theta d\theta$

~~$\frac{1}{x}$~~   $= \frac{1}{\tan \theta} = \frac{1}{\sin \theta} \cos \theta = \frac{\cos \theta}{\sin \theta}$

$$\int e^\theta \left( \frac{1 + \tan \theta + \tan^2 \theta}{1 + \tan^2 \theta} \right) \cdot \sec^2 \theta d\theta$$

$$= \int e^\theta (\sec^2 \theta + \tan \theta) d\theta = e^\theta \tan \theta + 1$$

$\uparrow$                        $\uparrow$   
 $f'$                        $f$

$$\ln^2 n = (\ln n)^2$$

Que:  $\int \frac{\ln n + 1}{(1 + \ln n)^2} dn = \int \frac{1}{1 + \ln n} - \frac{1}{(1 + \ln n)^2} dn$

$f(n) = \frac{1}{1 + \ln n}$   
 $f'(n) = -1 \times \frac{1}{(1 + \ln n)^2}$

$$= \int \left( \frac{1}{1 + \ln n} + n \cdot \frac{-1}{n(1 + \ln n)^2} \right) dn$$

$\frac{1}{y}$        $n$        $\frac{-1}{n}$

$$= n \cdot \frac{1}{1 + \ln n} + 1$$

Q2)  $\int \left( \ln(\ln n) + \frac{1}{\ln^2 n} \right) dn$

$f(y) = \ln y$

$f'(y) = \frac{1}{y}$  ;  $f''(y) = -\frac{1}{y^2}$

$\ln n = y$

$n = e^y$

$dn = e^y dy$

$\ln^2 n = (\ln n)^2$

Ans)  $\Rightarrow \int \left( \ln y + \frac{1}{y^2} \right) e^y dy = \int \left( \ln y + \frac{1}{y} - \frac{1}{y} + \frac{1}{y^2} \right) e^y$

$= \int \left( \ln y + \frac{1}{y} \right) e^y dy - \int \left( \frac{1}{y} + \left( -\frac{1}{y^2} \right) \right) e^y dy$

$= e^y \cdot \ln y - e^y \cdot \frac{1}{y} + 1$

$= e^y \left( \ln y - \frac{1}{y} \right) + 1 = n \left( \ln(\ln n) - \frac{1}{\ln n} \right) + 1$

Imp. ~~Imp.~~

Put  $n = \frac{1}{t}$  always necessary



रूट रूट (साथ वट्ट के साथ)

\* \* Metriculating <sup>→ Calculating</sup> Integration!

$$\begin{aligned}
 * \int \frac{dn}{n(n^6+1)} &= \int \frac{1}{n^7 \left(1 + \frac{1}{n^6}\right)} du \\
 &= -\frac{1}{6} \int \frac{dk}{1+k} = -\frac{1}{6} \ln(1+k) \quad \left[ \begin{array}{l} \frac{1}{n^6} = k \\ n^{-6} = k \\ -6 \frac{1}{n^7} dn = dk \end{array} \right.
 \end{aligned}$$

Method-2,

Put  $n = \frac{1}{t}$

$dn = -\frac{1}{t^2}$

$$\int \frac{-\frac{1}{t^2} dt}{\frac{1}{t} \left(\frac{1}{t^6} + 1\right)} = \int \frac{1}{t} \frac{t^6}{(1+t^6)} dt$$

$$= -\int \frac{t^5}{1+t^6} dt$$

$$= 1+t^6 = 1$$

Que:  $\int \frac{n^7}{(1-n^2)^5} dn = \int \frac{n^7}{n^{10} \left(\frac{1}{n^2} - 1\right)} dn$   $n \Rightarrow k = \frac{1}{n^2}$

$$= \frac{1}{n^3}$$

$$\frac{1}{\left(\frac{1}{n^2} - 1\right)^5} dn$$

$$\frac{1}{n^2} - 1 = t$$

$$-\frac{2}{n^3} dn = dt$$

Ques 1:  $\int \frac{x dx}{(1-x^4)^{3/2}}$

$$\int \frac{x dx}{x^6 \left(\frac{1}{x^4} + 1\right)^{3/2}}$$

$$= \int \frac{\frac{1}{x^5} dx}{\left(\frac{1}{x^4} + 1\right)^{3/2}}$$

$$x \frac{1}{x^4} = t$$

$$\frac{-4}{x^5} dx = dt$$

$$\frac{1}{x^4} + 1 = t$$

Ques 2:  $\int \frac{x-1}{x^2 \sqrt{2x^2-2x+1}} = \int \frac{\frac{1}{x^2} - \frac{1}{x^3}}{\sqrt{2 - \frac{2}{x} + \frac{1}{x^2}}}$

$$= 2 - \frac{2}{x} + \frac{1}{x^2} = t^2$$

2)  $\int \frac{dx}{x^4 (x^3+1)^2}$

$$\int \frac{dx}{x^{10} \left(1 + \frac{1}{x^3}\right)^2}$$

$$1 + \frac{1}{x^3} = t$$

$$Q. \int \frac{n^4 - 1}{n^2 \sqrt{n^4 + n^2 + 1}} dn = \int \frac{1 - \frac{1}{n^4}}{\sqrt{1 + \frac{1}{n^2} + \frac{1}{n^4}}} \frac{dn}{n}$$

$$= \int \frac{n^3 (n - \frac{1}{n^3}) dn}{n^3 \sqrt{n^2 + 1 + \frac{1}{n^2}}}$$

$$n^2 + 1 + \frac{1}{n^2} = k^2$$

$$\int \frac{k dk}{k}$$

$$(2n - \frac{2}{n^3}) dn = 2k dk$$

Que:  $\int \frac{(an^2 - b)}{n \sqrt{c^2 n^2 - (an^2 + b)^2}} dn$

$$= \int \frac{n^2 (a - \frac{b}{n^2}) dn}{n^2 \sqrt{c^2 - (an + \frac{b}{n})^2}}$$

$$an + \frac{b}{n} = t$$

$$(a - \frac{b}{n^2}) dn = dt$$

$$= \int \frac{dt}{\sqrt{c^2 - t^2}}$$

29/06/17

(8-1)

(25)

diff. both side

$$\frac{1 - (\cot n)^{2009}}{\sin n + (\cot n)^{2009}} = \frac{1}{k} \cdot \frac{1}{\left( \sin^n k_n + \cos^n k_n \right)^{k-1}} \left( k \sin^{k-1} n \cdot (-\cot n) \right)$$

$$= \frac{1}{\sin n^{2008}} \left( \sin^{2008} - \cos^{2008} \right)$$

$$\frac{\sin n}{\cos n} + \left( \frac{\cos n}{\sin n} \right)^{2009}$$

$$= \sin n \cos n \left( \frac{\sin^{k-2} n - \cos^{k-2} n}{\sin^k n + \cos^k n} \right)$$

$$\sin n \cos n \left( \frac{\sin^{2008} - \cos^{2008}}{\sin^{2010} + \cos^{2010}} \right)$$

$$= \frac{1}{\sin^{2008}} \left( \sin^{2008} - \cos^{2008} \right) \frac{\sin^{2010} + \cos^{2010}}{\sin^{2009} \cdot \cos n}$$

$$k = 2010$$

\* algebraic / trigonometric De Moivre's!

$$\int \frac{1}{x^4+1}$$

$$\int \sqrt{1-u^2} du$$

$$\int \frac{x^2}{x^4+1}$$

$$\int \sqrt{\cot u} du$$

$$\int \frac{1}{x^4+kx^2+1}$$

$$\int \frac{1}{\sin^4 u + \cos^4 u} du$$

$$\int \frac{m^2}{x^4+kx^2+1} dx$$

$$\int \frac{1}{\sin^6 u + \cos^6 u} du$$

$$\int \frac{\pm \sin u \pm \cos u}{1 + \sin u \cos u} du$$

Q.  $I = \int \frac{1}{x^4+1} dx = \int \frac{(1+x^2) + (1-x^2)}{x^4+1}$

$$= \int \frac{1+x^2}{1+x^4} + \int \frac{1-x^2}{1+x^4} = \int \frac{1 + \frac{1}{x^2}}{x^2 + \frac{1}{x^2}} dx = \int \frac{1 - \frac{1}{x^2}}{x^2 + \frac{1}{x^2}}$$

$$= \int \frac{dt}{t^2+2} - \int \frac{dp}{p^2-2}$$

$$\begin{aligned} \frac{x-1}{x} &= t \\ \left(1 + \frac{1}{x^2}\right) dx &= dt \end{aligned}$$

$$x^2 + \frac{1}{x^2} = t^2 + 2$$

Q. ~~Q. =~~

$$= x + \frac{1}{x} = p$$

$$\left(1 - \frac{1}{x^2}\right) dx = dp$$

$$x^2 + \frac{1}{x^2} + 2 = p^2$$



$$\textcircled{2} \quad I = \int \frac{x^2}{x^4+1} = \frac{1}{2} \int \frac{2x^2}{x^4+1} = \frac{1}{2}$$

$$\frac{1}{2} \int \frac{(x^2+1) + (x^2-1)}{x^4+1}$$

$$= \frac{1}{2} \int \frac{x^2+1}{x^4+1} dx + \frac{1}{2} \int \frac{x^2-1}{x^4+1} dx$$

~~$$\frac{1}{2} \int \frac{x^2+1}{x^4+1} dx + \frac{1}{2} \int \frac{x^2-1}{x^4+1} dx$$~~

$$\frac{1 + \frac{1}{x^2} = t}{-2}$$

$$2. \int \frac{dx}{x^2+px^2+1}$$

Sol:

Ques:  $\int \frac{2x^2}{x^4+1} dx$

(2)  $\int \frac{x^2+1}{x^4+3x^2+1} dx = \int \frac{1 + \frac{1}{x^2}}{x^2 + 3 + \frac{1}{x^2}}$   
 $= \int \frac{dt}{t^2+5}$

~~Ques:~~ Trigonometric twins

$\int \sqrt{\tan \theta} d\theta$

$\int p \cdot \frac{2k dk}{1+k^4}$

$= \int \frac{2k^2 dk}{1+k^4}$

$= \int \frac{(k^2+1) + (k^2-1)}{k^4+1} dk$

$\tan \theta = k^2$   
 $\sec^2 \theta d\theta = 2k dk$

$d\theta = \frac{2k dk}{\sec^2 \theta}$   
 $= \frac{2k dk}{1+\tan^2 \theta}$

$= \frac{2k dk}{1+k^4}$

Ques:  $\int \frac{d\theta}{\sin^4 \theta + \cos^4 \theta}$

$\rightarrow \tan \theta = k$   
 $\sec^2 \theta d\theta = dk$

$\int \frac{d\theta}{\cos^4 \theta (\tan^4 \theta + 1)} = \int \frac{\sec^4 \theta d\theta}{\tan^4 \theta + 1}$

$= \int \frac{(1+\tan^2 \theta)}{1+k^4} dk = \int \frac{1+k^2}{1+k^4} dk$   $\left( k - \frac{1}{k} = u \right)$   
 $= \int \frac{1 + \frac{1}{k^2}}{k^2 + \frac{1}{k^2}} dk$

$$Q \quad I = \int (\sqrt{\tan \theta} + \sqrt{\cot \theta}) d\theta$$

$$\Rightarrow \int \frac{\sin \theta}{\cos \theta} + \int \frac{\cos \theta}{\sin \theta}$$

$$= \int \frac{\sin \theta + \cos \theta}{\sin \theta \cos \theta} d\theta = \sqrt{2} \int \frac{dt}{\sqrt{1-t^2}} \quad \begin{matrix} \sin \theta = \cos \theta = t \\ (\cos \theta + \sin \theta) d\theta = dt \\ \sin^2 \theta + \cos^2 \theta = 2 \sin \theta \cos \theta \\ \frac{1-t^2}{2} = \sin \theta \cos \theta \end{matrix}$$

Ques:  $\int \frac{d\theta}{\sin^6 \theta + \cos^6 \theta}$

$$a^3 + b^3 = (a+b)(a^2 + b^2 - ab)$$

$$= \int \frac{\sec^4 \theta \sec^2 \theta}{1 + \tan^6 \theta} d\theta$$

$$= \int \frac{(1+t^2)^2}{1+t^6} dt$$

$$= \int \frac{(1+t^2)^2}{(1+t^2)(1+t^4-t^2)} dt$$

$$= \int \frac{1 + \frac{1}{t^2}}{t^2 + \frac{1}{t} - 1} dt \quad \text{Ans}$$

$$* \int \frac{\pm \sin n\theta + \cos m\theta}{1 + \sin n\theta \cos m\theta} d\theta$$

Ans

$$\int \frac{(\delta + c) + (\delta - c)}{1 + \delta c}$$

$$= \int \frac{\sin u + \cos u}{1 + \sin u \cos u} du + \int \frac{s - c}{1 + sc} du$$

$$= \int \frac{dt}{1 + \frac{1-t^2}{2}} + \int \frac{-dd}{1 + \frac{d^2-1}{2}}$$

$$\frac{\sin u + \cos u}{(\cos u - \sin u) du} = dd$$

$$\frac{\sin u - \cos u + 2}{(1 + \sin u + \cos u) du} = 2 \int \frac{dt}{3-t^2} - 2 \int \frac{dd}{1+d^2}$$

$$\sin^2 u + \cos^2 u + 2 \sin u \cos u = 1 + 2 \sin u \cos u$$

$$s^2 + c^2 + 2sc = 1 + 2sc$$

$$sc = \frac{d^2 - 1}{2}$$

$$1 - t^2 = 2sc$$

$$sc = \frac{1-t^2}{2}$$

$\int \rightarrow$

Ques)  $\int \frac{2 \cos u}{10 + \sin^2 u} du$

$$= \frac{\cos u + \sin u + \cos - \sin}{10 + 2 \sin u \cos u}$$

$$I = \int \frac{c + s}{10 + 2 \sin u \cos u} du + \int \frac{\cos - \sin}{10 + 2 \sin u \cos u}$$

$$= \int \frac{dt}{10 + t^2} + \int \frac{dd}{10 + d^2}$$

V. Pimp.

H.W ⇒ all full comp

$$\left(n \pm \frac{4}{n}\right)^2 = n^2 + \frac{16}{n^2} \pm 8$$

$$Q. \int \frac{e^x dx}{x^4 + 16} = \int \frac{dx}{x^2 \left(x^2 + \frac{16}{x^2}\right)}$$

$$\left(n \pm \frac{4}{n}\right)^2 =$$

$$\left(n \pm \frac{4}{n}\right)^2 = n^2 + \frac{16}{n^2} \pm 8$$

$$\frac{1}{8} \int \frac{e^x dx}{x^2 \left(x^2 + \frac{16}{x^2}\right)}$$

$$= \frac{1}{8} \int \frac{\left(\frac{4}{x^2} + 1\right) + \left(\frac{4}{x^2} - 1\right)}{x^2 + \frac{16}{x^2}}$$

$$\begin{aligned} n + \frac{4}{n} &= t \\ \left(1 - \frac{4}{x^2}\right) dx &= dt \\ \left(1 + \frac{4}{x^2}\right) dx &= dt \end{aligned}$$

$$x^2 + \frac{16}{x^2} = t^2 - 8$$

$$x^2 + \frac{16}{x^2} = t^2 + 8$$

$$= \frac{1}{8} \int \frac{1 + \frac{4}{x^2}}{x^2 + \frac{16}{x^2}} dx - \frac{1}{8} \int \frac{1 - \frac{4}{x^2}}{x^2 + \frac{16}{x^2}} dx$$

$$= \frac{1}{8} \int \frac{dt}{t^2 - 8} - \frac{1}{8} \int \frac{dy}{y^2 + 8}$$

Ques!  $\int \frac{x^2 + 3}{x^4 + 19x^2 + 19x} dx = \frac{1 + \frac{3}{x^2}}{x^2 + 19 + \frac{9}{x^2}}$

$$= \int \frac{1 + \frac{3}{x^2}}{x^2 + 19 + \frac{9}{x^2}}$$

$$\begin{aligned} x - \frac{3}{x} &= t \\ \int \left(1 + \frac{3}{x^2}\right) dx &= dt \end{aligned}$$

$$= \int \frac{dt}{t^2 + 25}$$

$$x^2 + \frac{9}{x^2} = t^2 + 6$$

Imp

\*  $\int \frac{1}{A\sqrt{B}}$  Integration of the form

where A and B are linear or quadratic expression

L - Linear  
Q - Quadratic

| A | B | Substitution.           | Result (into Linear)                                               |
|---|---|-------------------------|--------------------------------------------------------------------|
| L | L | → Put $B = t^2$         | $(an^2+bn+c) = a(n-\alpha)(n-\beta)$                               |
| L | L | → Put $B = t^2$         | (1) $\frac{1}{an^2+bn+c} = \frac{A}{n-\alpha} + \frac{B}{n-\beta}$ |
| L | Q | → Put $A = \frac{1}{t}$ | $\therefore I = A \int \frac{dn}{(n-\alpha)\sqrt{pn^2+qn+r}}$      |
| Q | Q |                         | (1) $+ B \int \frac{dn}{(n-\beta)\sqrt{pn^2+qn+r}}$                |

goes to twin

$(an^2+bn+c) \int \frac{dn}{pn^2+qn+r}$

a

$$\int \frac{1}{(n-1)\sqrt{n-2}}$$

A = (n-1)  
B = (n-2)

a

$$\int \frac{dn}{(n^2+n+1)\sqrt{n+1}}$$

L      L

(2) if  $an^2+bn+c = (ln+m)^2$

$$\therefore I = \int \frac{dn}{(ln+m)^2 \sqrt{pn^2+qn+r}}$$

$$ln+m = \frac{1}{t}$$

b = q = 0 Imp

$$I = \int \frac{dn}{(a^2+c)\sqrt{pn^2+r}}$$

Put  $n = \frac{1}{t}$

$$\text{Ques 1} \int \frac{dn}{(n-1)\sqrt{n-2}}$$

$$= \begin{aligned} A &= (n-1) \\ B &= (n-2) \end{aligned}$$

$$\begin{aligned} &\frac{2+dt}{(n-1)\sqrt{2+dt}} \\ &2 \int \frac{dt}{t^2+1} \end{aligned}$$

$$\begin{aligned} &= \int \frac{dn}{(n-1)\sqrt{t^2}} \\ (n-2) &= t^2 \\ |dn &= 2t dt \end{aligned}$$

Q.2

$$\int \frac{dn}{(n-1)\sqrt{n^2+n+1}}$$

$$= -\frac{1}{t} \int \frac{dt}{\sqrt{3t^2+3t+1}}$$

$$= \int \frac{dt}{\sqrt{3t^2+3t+1}}$$

$$n-1 = \frac{1}{t}$$

$$n+1 = \frac{1}{t}$$

$$dn = -\frac{1}{t^2} dt$$

$$n = \frac{1}{t} + 1 = \frac{t+1}{t}$$

$$n^2+n+1 = \left(\frac{t+1}{t}\right)^2 + \frac{t+1}{t} + 1$$

$$= \frac{(t+1)^2 + t(t+1) + t^2}{t^2}$$

$$= \frac{3t^2+3t+1}{t^2}$$

$$\text{Q} \int \frac{dn}{(n^2-2n+1)\sqrt{n^2+nt+1}}$$

$$= \int \frac{dn}{(n-1)^2 \sqrt{n^2+nt+1}} = \int \frac{t}{\sqrt{3t^2+3t+1}} dt$$

Q.

$$\int \frac{8 \, dn}{(n^2 + 5n + 2) \sqrt{n-2}}$$

$$n-2 = t^2 \\ dn = 2t \, dt$$

$$\rightarrow \int \frac{8t \, dt}{n^2 + 5n + 2 \times t}$$

$$\rightarrow \int \frac{8 \, dt}{t^2 + at^2 + 16}$$

$$t = \frac{4}{t}$$

$$= \int \frac{\frac{8}{t^2} \, dt}{t^2 + \left(\frac{4}{t}\right)^2 + 9}$$

$$t = \frac{4}{t^2}$$

$$= \int \frac{\left(1 + \frac{4}{t^2}\right) - \left(1 - \frac{4}{t^2}\right) \, du}{t^2 + \left(\frac{4}{t}\right)^2 + 9}$$

$$= \int \frac{1 + \frac{4}{t^2}}{t^2 + \left(\frac{4}{t}\right)^2 + 9} \, dt = \int \frac{\left(1 - \frac{4}{t^2}\right)}{t^2 + \left(\frac{4}{t}\right)^2 + 9} \, dt$$

$$t - \frac{4}{t} = \lambda$$

$$t + \frac{4}{t} = \rho$$

Ques:

$$\int \frac{dn}{(n^2 - n) \sqrt{n^2 + n + 1}}$$

$$\frac{1}{n^2 - 1} = \frac{1}{n-1} - \frac{1}{n}$$

$$\int \frac{dn}{(n-1) \sqrt{n^2 + n + 1}} - \int \frac{dn}{n \sqrt{n^2 + n + 1}}$$

$$\uparrow n = \frac{1}{t}$$



$$Q. \int \frac{dn}{(n^2+3n+2)\sqrt{n-1}}$$

$$= \int \frac{1}{n^2-3n+2} - \frac{1}{(n-1)(n-2)}$$

$$\sqrt{n-1}$$

$$n-1 = t^2$$

$$dn = 2t dt$$

$$= \frac{1}{n-2} - \frac{1}{n-1}$$

$$= \int \frac{1}{(n+2)\sqrt{n+1}} dn - \int \frac{dn}{(n-1)\sqrt{n+1}}$$

Sol:

$$= \int \frac{f du}{u^2}$$

$$n = t^2 - 1$$

$$n+1 = t^2$$

$$dn = 2t dt$$

$$= \int \frac{2t dt}{t^2}$$

$$= \int \frac{2 dt}{t}$$

Que:

$$\int \frac{dn}{(n^2+1)\sqrt{n^2-2}}$$

$$n = \frac{1}{t}$$

$$dn = -\frac{1}{t^2} dt$$

$$-t dt = \frac{1}{2} u du$$

$$1-2t^2 = u^2$$

$$-4t dt = 2u du$$

$$-2t dt = u du$$

$$t^2 = \frac{1-u^2}{2}$$

$$1+t^2 = 1 + \frac{1-u^2}{2}$$

$$= - \int \frac{t dt}{(1+t^2)(\sqrt{1-2t^2})}$$

$$= \frac{1}{2} \int \frac{u du}{u(3-u^2)}$$

$$= \int \frac{du}{3-u^2}$$

A

$$n=2 \quad f = \sin \theta$$

\* Reduction formula!

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$I_n = \int \sin^n \theta \, d\theta$$

$$= \frac{\sin^n}{II} \cdot \frac{\sin^{n-1}}{I} \, d\theta$$

Break  $\sin^{n-1}$   
 into part line

$$= -\cos \theta \sin^{n-1} - (n-1) \int (-\cos \theta) \cdot \sin^{n-2} \cdot \cos \theta \, d\theta$$

$$= -\cos \theta \cdot \sin^{n-1} - (n-1) \int \sin^2 \theta \cdot \sin^{n-2} \, d\theta$$

$$I_n = -\cos \theta (\sin \theta)^{n-1} - (n-1) \int \sin^2 \theta \, d\theta + (n-1) \int \sin^{n-2} \theta \, d\theta$$

$$= -\cos \theta (\sin \theta)^{n-1} - (n-1) I_n + (n-1) I_{n-2}$$

$$(n-1+1) I_n = -\cos \theta \sin^{n-1} + (n-1) I_{n-2}$$

$$I_n = \frac{-1}{n} \cdot \cos \theta \sin^{n-1} + \frac{n-1}{n} I_{n-2}$$

$$I_3 = \frac{-1}{3} \cos \theta \sin^2 \theta + \frac{2}{3} I_1$$

$$I_3 = \int \sin^3 \theta \, d\theta$$

$$I_1 = \int \sin \theta \, d\theta = -\cos \theta$$

Learn

Sub

$$Q \quad I_n = \int \tan^n x \, dx$$

$$\int \tan^2 x - \tan^{n-2} x \, dx$$

$$I_3 = \int \sin^3 x \, dx$$

$$I_1 = \int \sin x \, dx = -\cos x$$

$$\int (\sec^2 x - 1) \tan^{n-2} x \, dx$$

$$I_n = \int \sec^2 x \cdot \tan^{n-2} x \, dx - \int \tan^{n-2} x \, dx$$

$$I_n + I_{n-2} = \int \sec^2 x \cdot \tan^{n-2} x \, dx$$

$$= \int t^{n-2} \, dt = \frac{t^{n-1}}{n-1} = \frac{(\tan x)^{n-1}}{n-1}$$

Q2

$$I_n = \int \sec^n x \, dx$$

$$= \int \underbrace{\sec^2 x}_I \cdot \underbrace{\sec^{n-2} x}_{II} \, dx$$

$$= \sec^{n-2} x \tan x - (n-2) \int \sec^{n-3} x \cdot \sec x \tan x \cdot \tan x \, dx$$

$$= \sec^{n-2} x \cdot \tan x - (n-2) \int \sec^{n-2} x \cdot (\sec^2 x - 1) \, dx$$

$$= \sec^{n-2} x \tan x - (n-2) \int \sec^n x \, dx + (n-2) \int \sec^{n-2} x \, dx$$

$$\boxed{I_n = \sec^{n-2} x \tan x - (n-2) I_n + (n-2) I_{n-2}}$$

17/17

LATE

⇒ Full complete

$$Q. I_n = \int \frac{dx}{(x^2+a^2)^n}$$

$$\int \frac{1 \cdot dx}{(x^2+a^2)^n} =$$

$$n \frac{1}{(x^2+a^2)^n} - \left( \frac{(-n) \cdot (2x)}{(x^2+a^2)^{n+1}} \right) x \cdot dx$$

$$= \frac{n}{(x^2+a^2)^n} + 2n \int \frac{(x^2+a^2) - a^2}{(x^2+a^2)^{n+1}} dx$$

$$I_n = \frac{n}{(x^2+a^2)^n} + 2n \int \frac{dx}{(x^2+a^2)^n} - 2na^2 \int \frac{dx}{(x^2+a^2)^{n+1}}$$

$$I_n = \frac{n}{(x^2+a^2)^n} + 2n I_n - 2na^2 I_{n+1}$$

# SBG STUDY