

Solved Paper 2015





JEE Main Joint Entrance Examination

Instructions

- This test consists of 90 questions.
- There are three parts in the question paper A, B, C consisting of Physics, Chemistry & Mathematics having 30 questions in each part of equal weightage. Each question is allotted 4 marks for correct response.
- Candidates will be awarded marks as stated above in instruction no. 2 for correct response of each question. 1 marks will be deducted for indicating incorrect response of each question. No deduction from the total score will be made if no response is indicated for an item in the answer sheet.
- There is only one correct response for each question. Filling up more than one response in any question will be treated as wrong response and marks for wrong response will be deducted according as per instructions.


Physics

1. Two stones are thrown up simultaneously from the edge of a cliff 240 m high with initial speed of 10 m/s and 40 m/s respectively. Which of the following graph best represents the time variation of relative position of the second stone with respect to the first? Assume stones do not rebound after hitting the ground and neglect air resistance, take $g = 10 \text{ m/s}^2$



2. The period of oscillation of a simple pendulum is $T = 2\pi \sqrt{\frac{L}{g}}$. Measured value of L is 20.0 cm known to 1 mm accuracy and time for 100 oscillations of the pendulum is found to be 90 s using a wrist watch of 1s resolution. The accuracy in the determination of g is
- (a) 2% (b) 3%
(c) 1% (d) 5%

3. Given in the figure are two blocks A and B of weight 20 N and 100 N respectively. These are being pressed against a wall by a force F as shown in figure. If the coefficient of friction between the blocks is 0.1 and between block B and the wall is 0.15, the frictional force applied by the wall in block B is



- (a) 100 N (b) 80 N
(c) 120 N (d) 150 N

2 JEE Main Solved Paper 2015

4. A particle of mass m moving in the x -direction with speed $2v$ is hit by another particle of mass $2m$ moving in the y -direction with speed v . If the collision is perfectly inelastic, the percentage loss in the energy during the collision is close to

- (a) 44% (b) 50%
(c) 56% (d) 62%


5. Distance of the centre of mass of a solid uniform cone from its vertex is z_0 . If the radius of its base is R and its height is h , then z_0 is equal to

- (a) $\frac{h^2}{4R}$ (b) $\frac{3h}{4}$
(c) $\frac{5h}{8}$ (d) $\frac{3h^2}{8R}$

6. From a solid sphere of mass M and radius R , a cube of maximum possible volume is cut. Moment of inertia of cube about an axis passing through its centre and perpendicular to one of its faces is

- (a) $\frac{MR^2}{32\sqrt{2}\pi}$ (b) $\frac{MR^2}{16\sqrt{2}\pi}$
(c) $\frac{4MR^2}{9\sqrt{3}\pi}$ (d) $\frac{4MR^2}{3\sqrt{3}\pi}$

7. From a solid sphere of mass M and radius R , a spherical portion of radius $\left(\frac{R}{2}\right)$ is removed as shown in the figure. Taking gravitational potential $V = 0$ at $r = \infty$, the potential at the centre of the cavity thus formed is ($G =$ gravitational constant)



- (a) $-\frac{GM}{2R}$
(b) $-\frac{GM}{R}$
(c) $-\frac{2GM}{3R}$
(d) $-\frac{2GM}{R}$

8. A pendulum made of a uniform wire of cross-sectional area A has time period T . When an additional mass M is added to its bob, the time period changes T_M . If the Young's modulus of the material of the wire is Y , then $\frac{1}{Y}$ is equal to ($g =$ gravitational acceleration)

- (a) $\left[\left(\frac{T_M}{T}\right)^2 - 1\right] \frac{A}{Mg}$
(b) $\left[\left(\frac{T_M}{T}\right)^2 - 1\right] \frac{Mg}{A}$
(c) $\left[1 - \left(\frac{T_M}{T}\right)^2\right] \frac{A}{Mg}$
(d) $\left[1 - \left(\frac{T}{T_M}\right)^2\right] \frac{A}{Mg}$

9. Consider a spherical shell of radius R at temperature T . The black body radiation inside it can be considered as an ideal gas of photons with internal energy per unit volume $u = \frac{U}{V} \propto T^4$ and pressure $p = \frac{1}{3} \left(\frac{U}{V}\right)$. If the shell now undergoes an adiabatic expansion, the relation between T and R is

- (a) $T \propto e^{-R}$ (b) $T \propto e^{-3R}$
(c) $T \propto \frac{1}{R}$ (d) $T \propto \frac{1}{R^3}$

10. A solid body of constant heat capacity $1 \text{ J}^\circ\text{C}$ is being heated by keeping it in contact with reservoirs in two ways

- (i) Sequentially keeping in contact with 2 reservoirs such that each reservoir supplies same amount of heat.
(ii) Sequentially keeping in contact with 8 reservoirs such that each reservoir supplies same amount of heat.

In both the cases, body is brought from initial temperature 100°C to final temperature 200°C . Entropy change of the body in the two cases respectively, is


- (a) $\ln 2, 4\ln 2$
(b) $\ln 2, \ln 2$
(c) $\ln 2, 2\ln 2$
(d) $2\ln 2, 8\ln 2$

11. Consider an ideal gas confined in an isolated closed chamber. As the gas undergoes an adiabatic expansion, the average time of collision between molecules increases as V^q , where V is the volume of the gas. The

value of q is $\left(\gamma = \frac{C_p}{C_v}\right)$

- (a) $\frac{3\gamma + 5}{6}$ (b) $\frac{3\gamma - 5}{6}$ (c) $\frac{\gamma + 1}{2}$ (d) $\frac{\gamma - 1}{2}$


12. For a simple pendulum, a graph is plotted between its Kinetic Energy (KE) and Potential Energy (PE) against its displacement d . Which one of the following represents these correctly? (graphs are schematic and not drawn to scale)



13. A train is moving on a straight track with speed 20 ms^{-1} . It is blowing its whistle at the frequency of 1000 Hz . The percentage change in the frequency heard by a person standing near the track as the train passes him is close to (speed of sound = 320 ms^{-1})

- (a) 6% (b) 12% (c) 18% (d) 24%



14. A long cylindrical shell carries positive surface charge σ in the upper half and negative surface charge $-\sigma$ in the lower half. The electric field lines around the cylinder will look like figure given in (figures are schematic and not drawn to scale)



15. A uniformly charged solid sphere of radius R has potential V_0 (measured with respect to ∞) on its surface. For this sphere, the equipotential surfaces with potentials $\frac{3V_0}{2}$, $\frac{5V_0}{4}$, $\frac{3V_0}{4}$ and $\frac{V_0}{4}$ have radius R_1 , R_2 , R_3 , and R_4 respectively. Then,

- (a) $R_1 = 0$ and $R_2 > (R_4 - R_3)$
 (b) $R_1 \neq 0$ and $(R_2 - R_1) > (R_4 - R_3)$
 (c) $R_1 = 0$ and $R_2 < (R_4 - R_3)$
 (d) $2R < R_4$

16. In the given circuit, charge Q_2 on the $2 \mu\text{F}$ capacitor changes as C is varied from $1 \mu\text{F}$ to $3 \mu\text{F}$. Q_2 as a function of C is given properly by (figures are drawn schematically and are not to scale)




17. When 5 V potential difference is applied across a wire of length 0.1 m , the drift speed of electrons is $2.5 \times 10^{-4} \text{ ms}^{-1}$. If the electron density in the wire is $8 \times 10^{28} \text{ m}^{-3}$ the resistivity of the material is close to

- (a) $1.6 \times 10^{-8} \Omega\text{m}$
 (b) $1.6 \times 10^{-7} \Omega\text{m}$
 (c) $1.6 \times 10^{-6} \Omega\text{m}$
 (d) $1.6 \times 10^{-5} \Omega\text{m}$

4 JEE Main Solved Paper 2015

18. In the circuit shown below, the current in the 1Ω resistor is




- (a) 1.3 A, from P to Q
- (b) 0 A
- (c) 0.13 A, from Q to P
- (d) 0.13 A, from P to Q

19. Two coaxial solenoids of different radii carry current I in the same direction. Let F_1 be the magnetic force on the inner solenoid due to the outer one and F_2 be the magnetic force on the outer solenoid due to the inner one. Then,


- (a) $F_1 = F_2 = 0$
- (b) F_1 is radially inwards and F_2 is radially outwards
- (c) F_1 is radially inwards and $F_2 = 0$
- (d) F_1 is radially outwards and $F_2 = 0$

20. Two long current carrying thin wires, both with current I , are held by insulating threads of length L and are in equilibrium as shown in the figure, with threads making an angle θ with the vertical. If wires have mass λ per unit length then, the value of I is ($g =$ gravitational acceleration)



- (a) $\sin\theta \sqrt{\frac{\pi\lambda gL}{\mu_0 \cos\theta}}$
- (b) $2\sin\theta \sqrt{\frac{\pi\lambda gL}{\mu_0 \cos\theta}}$
- (c) $2 \sqrt{\frac{\pi gL}{\mu_0} \tan\theta}$
- (d) $\sqrt{\frac{\pi\lambda gL}{\mu_0} \tan\theta}$


21. A rectangular loop of sides 10 cm and 5 cm carrying a current I of 12 A is placed in different orientations as shown in the figures below.



If there is a uniform magnetic field of 0.3 T in the positive z -direction in which orientations the loop would be in (i) stable equilibrium and (ii) unstable equilibrium?

- (a) (a) and (b) respectively
- (b) (a) and (c) respectively
- (c) (b) and (d) respectively
- (d) (b) and (c) respectively

22. An inductor ($L = 0.03$ H) and a resistor ($R = 0.15$ k Ω) are connected in series to a battery of 15V EMF in a circuit shown below. The key K_1 has been kept closed for a long time. Then at $t = 0$, K_1 is opened and key K_2 is closed simultaneously. At $t = 1$ ms, the current in the circuit will be ($e^5 \cong 150$)




- (a) 100 mA
- (b) 67 mA
- (c) 6.7 mA
- (d) 0.67 mA

23. A red LED emits light at 0.1 W uniformly around it. The amplitude of the electric field of the light at a distance of 1 m from the diode is

- (a) 1.73 V/m
- (b) 2.45 V/m
- (c) 5.48 V/m
- (d) 7.75 V/m

24. Monochromatic light is incident on a glass prism of angle A . If the refractive index of the material of the prism is μ , a ray incident at an angle θ , on the face AB would get transmitted through the face AC of the prism provided



- (a) $\theta > \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$
 (b) $\theta < \sin^{-1} \left[\mu \sin \left(A - \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$
 (c) $\theta > \cos^{-1} \left[\mu \sin \left(A + \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$
 (d) $\theta < \cos^{-1} \left[\mu \sin \left(A + \sin^{-1} \left(\frac{1}{\mu} \right) \right) \right]$

25. On a hot summer night, the refractive index of air is smallest near the ground and increases with height from the ground. When a light beam is directed horizontally, the Huygens principle leads us to conclude that as it travels, the light beam

- (a) becomes narrower
 (b) goes horizontally without any deflection
 (c) bends downwards
 (d) bends upwards

26. Assuming human pupil to have a radius of 0.25 cm and a comfortable viewing distance of 25 cm, the minimum separation between two objects that human eye can resolve at 500 nm wavelength is

- (a) $1 \mu\text{m}$ (b) $30 \mu\text{m}$ (c) $100 \mu\text{m}$ (d) $300 \mu\text{m}$

27. As an electron makes a transition from an excited state to the ground state of a hydrogen like atom/ion

- (a) its kinetic energy increases but potential energy and total energy decrease
 (b) kinetic energy, potential energy and total energy decrease
 (c) kinetic energy decreases, potential energy increases but total energy remains same
 (d) kinetic energy and total energy decrease but potential energy increases

28. Match List-I (fundamental experiment) with List-II (its conclusion) and select the correct option from the choices given below the list.


List I	List II
A. Franck-Hertz experiment	1. Particle nature of light
B. Photo-electric experiment	2. Discrete energy levels of atom
C. Davisson-Germer experiment	3. Wave nature of electron
	4. Structure of atom

A	B	C	A	B	C
(a) 1	4	3	(b) 2	4	3
(c) 2	1	3	(d) 4	3	2


29. A signal of 5 kHz frequency is amplitude modulated on a carrier wave of frequency 2MHz. The frequencies of the resultant signal is/are

- (a) 2 MHz only
 (b) 2005 kHz and 1995 kHz
 (c) 2005 kHz 2000 kHz and 1995 kHz
 (d) 2000 kHz and 1995 kHz

30. An LCR circuit is equivalent to a damped pendulum. In an LCR circuit, the capacitor is charged to Q_0 and then connected to the L and R as shown below.



If a student plots graphs of the square of maximum charge (Q_{Max}^2) on the capacitor with time (t) for two different values L_1 and L_2 ($L_1 > L_2$) of L , then which of the following represents this graph correctly? (plots are schematic and not drawn to scale)



Chemistry

1. The molecular formula of a commercial resin used for exchanging ions in water softening is $C_8H_7SO_3Na$ (mol. wt. = 206). What would be the maximum uptake of Ca^{2+} ions by the resin when expressed in mole per gram resin?

- (a) $\frac{1}{103}$ (b) $\frac{1}{206}$
 (c) $\frac{2}{309}$ (d) $\frac{1}{412}$

2. Sodium metal crystallises in a body centred cubic lattice with a unit cell edge of 4.29 Å. The radius of sodium atom is approximately

- (a) 1.86 Å (b) 3.22 Å (c) 5.72 Å (d) 0.93 Å


3. Which of the following is the energy of a possible excited state of hydrogen?

- (a) +13.6 eV
 (b) -6.8 eV
 (c) -3.4 eV
 (d) +6.8 eV

4. The intermolecular interaction that is dependent on the inverse cube of distance between the molecules is

- (a) ion-ion interaction (b) ion-dipole interaction
 (c) London force (d) hydrogen bond

5. The following reaction is performed at 298K



The standard free energy of formation of NO (g) is 86.6 kJ/mol at 298 K. What is the standard free energy of formation of $NO_2(g)$ at 298 K? ($K_p = 1.6 \times 10^{12}$)

- (a) $R(298) \ln (1.6 \times 10^{12}) - 86600$
 (b) $86600 + R(298) \ln (1.6 \times 10^{12})$
 (c) $86600 - \frac{\ln (1.6 \times 10^{12})}{R(298)}$
 (d) $0.5 [2 \times 86600 - R(298) \ln (1.6 \times 10^{12})]$

6. The vapour pressure of acetone at 20°C is 185 torr. When 1.2 g of a non-volatile substance was dissolved in 100 g of acetone at 20°C, its vapour pressure was 183 torr. The molar mass ($g \text{ mol}^{-1}$) of the substance is

- (a) 32 (b) 64
 (c) 128 (d) 488

7. The standard Gibbs energy change at 300K for the reaction, $2A \rightleftharpoons B + C$ is 2494.2 J. At a given time, the composition of the reaction mixture is $[A] = \frac{1}{2}$, $[B] = 2$

and $[C] = \frac{1}{2}$. The reaction proceeds in the

$[R = 8.314 \text{ JK/mol}, e = 2.718]$

- (a) forward direction because $Q > K_c$
 (b) reverse direction because $Q > K_c$
 (c) forward direction because $Q < K_c$
 (d) reverse direction because $Q < K_c$

8. Two Faraday of electricity is passed through a solution of $CuSO_4$. The mass of copper deposited at the cathode is (at. mass of Cu = 63.5 u)

- (a) 0g (b) 63.5g (c) 2g (d) 127g

9. Higher order (>3) reactions are rare due to

- (a) low probability of simultaneous collision of all the reacting species
 (b) increase in entropy and activation energy as more molecules are involved
 (c) shifting of equilibrium towards reactants due to elastic collisions
 (d) loss of active species on collision

10. 3 g of activated charcoal was added to 50 mL of acetic acid solution (0.06 N) in a flask. After an hour it was filtered and the strength of the filtrate was found to be 0.042 N. The amount of acetic acid adsorbed (per gram of charcoal) is

- (a) 18 mg (b) 36 mg (c) 42 mg (d) 54 mg

11. The ionic radii (in Å) of N^{3-} , O^{2-} and F^- respectively are

- (a) 1.36, 1.40 and 1.71 (b) 1.36, 1.71 and 1.40
 (c) 1.71, 1.40 and 1.36 (d) 1.71, 1.36 and 1.40

12. In the context of the Hall-Heroult process for the extraction of Al, which of the following statements is false?

- (a) CO and CO_2 are produced in this process
 (b) Al_2O_3 is mixed with CaF_2 which lowers the melting point of the mixture and brings conductivity
 (c) Al^{3+} is reduced at the cathode to form Al
 (d) Na_3AlF_6 serves as the electrolyte

13. From the following statements regarding H_2O_2 , choose the incorrect statement.

- (a) It can act only as an oxidising agent
- (b) It decomposed on exposure to light
- (c) It has to be stored in plastic or wax lined glass bottles in dark
- (d) It has to be kept away from dust

14. Which one of the following alkaline earth metal sulphates has its hydration enthalpy greater than its lattice enthalpy?

- (a) $CaSO_4$
- (b) $BeSO_4$
- (c) $BaSO_4$
- (d) $SrSO_4$

15. Which among the following is the most reactive?

- (a) Cl_2
- (b) Br_2
- (c) I_2
- (d) ICl

16. Match the catalysts to the correct processes.

Catalyst	Process
(A) $TiCl_3$	(i) Wacker process
(B) $PdCl_2$	(ii) Ziegler-Natta polymerisation
(C) $CuCl_2$	(iii) Contact process
(D) V_2O_5	(iv) Deacon's process

- (a) (A) - (iii), (B) - (ii), (C) - (iv), (D) - (i)
- (b) (A) - (ii), (B) - (i), (C) - (iv), (D) - (iii)
- (c) (A) - (ii), (B) - (iii), (C) - (iv), (D) - (i)
- (d) (A) - (iii), (B) - (i), (C) - (ii), (D) - (iv)

17. Which one has the highest boiling point?

- (a) He
- (b) Ne
- (c) Kr
- (d) Xe

18. The number of geometric isomers that can exist for square planar $[Pt (Cl) (py) (NH_3) (NH_2OH)]^+$ is (py = pyridine)

- (a) 2
- (b) 3
- (c) 4
- (d) 6

19. The colour of $KMnO_4$ is due to

- (a) $M \rightarrow L$ charge transfer transition
- (b) $d-d$ transition
- (c) $L \rightarrow M$ charge transfer transition
- (d) $\sigma \rightarrow \sigma^*$ transition

20. Assertion (A) Nitrogen and oxygen are the main components in the atmosphere but these do not react to form oxides of nitrogen.

Reason (R) The reaction between nitrogen and oxygen requires high temperature.

- (a) Both Assertion and Reason are correct and the reason is the correct explanation for the Assertion.
- (b) Both Assertion and Reason are correct but the reason is not the correct explanation for the Assertion.
- (c) The Assertion is incorrect but the Reason is correct.
- (d) Both the Assertion and Reason are incorrect.

21. In Carius method of estimation of halogens, 250 mg of an organic compound gave 141 mg of $AgBr$. The percentage of bromine in the compound is


(at. mass $Ag = 108, Br = 80$)

- (a) 24
- (b) 36
- (c) 48
- (d) 60

22. Which of the following compound will exhibit geometrical isomerism?

- (a) 1-phenyl-2-butene
- (b) 3-phenyl-1-butene
- (c) 2-phenyl-1-butene
- (d) 1, 1-diphenyl-1-propane

23. Which compound would give 5-keto-2-methyl hexanal upon ozonolysis?




24. The synthesis of alkyl fluorides is best accomplished by

- (a) free radical fluorination
- (b) Sandmeyer's reaction
- (c) Finkelstein reaction
- (d) Swarts reaction

8 JEE Main Solved Paper 2015


25. In the following sequence of reaction,



The product C is

- (a) $\text{C}_6\text{H}_5\text{COOH}$ (b) $\text{C}_6\text{H}_5\text{CH}_3$
(c) $\text{C}_6\text{H}_5\text{CH}_2\text{OH}$ (d) $\text{C}_6\text{H}_5\text{CHO}$

26. In the reaction,



The product E is

- (a) (b) (c) (d)

27. Which polymer is used in the manufacture of paints and lacquers?

- (a) Bakelite
(b) Glyptal
(c) Polypropene
(d) Polyvinyl chloride

28. Which of the vitamins given below is water soluble?

- (a) Vitamin C
(b) Vitamin D
(c) Vitamin E
(d) Vitamin K

29. Which of the following compounds is not an antacid?

- (a) Aluminium hydroxide
(b) Cimetidine
(c) Phenelzine
(d) Ranitidine

30. Which of the following compounds is not coloured yellow?

- (a) $\text{Zn}_2[\text{Fe}(\text{CN})_6]$
(b) $\text{K}_3[\text{Co}(\text{NO}_2)_6]$
(c) $(\text{NH}_4)_3[\text{As}(\text{Mo}_3\text{O}_{10})_4]$
(d) BaCrO_4

Mathematics

1. Let A and B be two sets containing four and two elements respectively. Then, the number of subsets of the set $A \times B$, each having atleast three elements are

- (a) 219 (b) 256 (c) 275 (d) 510

2. A complex number z is said to be unimodular, if $|z| = 1$. Suppose z_1 and z_2 are

complex numbers such that $\frac{z_1 - 2z_2}{2 - z_1\bar{z}_2}$ is

unimodular and z_2 is not unimodular. Then, the point z_1 lies on a

- (a) straight line parallel to X-axis.
(b) straight line parallel to Y-axis.
(c) circle of radius 2.
(d) circle of radius $\sqrt{2}$.

3. Let α and β be the roots of equation $x^2 - 6x - 2 = 0$. If $a_n = \alpha^n - \beta^n$, for $n \geq 1$, then

the value of $\frac{a_{10} - 2a_8}{2a_9}$ is equal to

- (a) 6 (b) -6
(c) 3 (d) -3

4. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ a & 2 & b \end{bmatrix}$ is a matrix satisfying the

equation $AA^T = 9I$, where I is 3×3 identity matrix, then the ordered pair (a, b) is equal to

- (a) (2, -1) (b) (-2, 1)
(c) (2, 1) (d) (-2, -1)

5. The set of all values of λ for which the system of linear equations $2x_1 - 2x_2 + x_3 = \lambda x_1$, $2x_1 - 3x_2 + 2x_3 = \lambda x_2$ and $-x_1 + 2x_2 = \lambda x_3$ has a non-trivial solution,
 (a) is an empty set
 (b) is a singleton set
 (c) contains two elements
 (d) contains more than two elements

6. The number of integers greater than 6000 that can be formed, using the digits 3, 5, 6, 7 and 8 without repetition, is
 (a) 216 (b) 192
 (c) 120 (d) 72

7. The sum of coefficients of integral powers of x in the binomial expansion of $(1 - 2\sqrt{x})^{50}$ is
 (a) $\frac{1}{2}(3^{50} + 1)$ (b) $\frac{1}{2}(3^{50})$
 (c) $\frac{1}{2}(3^{50} - 1)$ (d) $\frac{1}{2}(2^{50} + 1)$

8. If m is the AM of two distinct real numbers l and n ($l, n > 1$) and G_1, G_2 and G_3 are three geometric means between l and n , then $G_1^4 + 2G_2^4 + G_3^4$ equals
 (a) $4l^2mn$ (b) $4lm^2n$
 (c) $4lmn^2$ (d) $4l^2m^2n^2$

9. The sum of first 9 terms of the series $\frac{1^3}{1} + \frac{1^3 + 2^3}{1 + 3} + \frac{1^3 + 2^3 + 3^3}{1 + 3 + 5} + \dots$ is
 (a) 71 (b) 96
 (c) 142 (d) 192

10. $\lim_{x \rightarrow 0} \frac{(1 - \cos 2x)(3 + \cos x)}{x \tan 4x}$ is equal to
 (a) 4 (b) 3
 (c) 2 (d) $\frac{1}{2}$

11. If the function $g(x) = \begin{cases} k\sqrt{x+1} & , 0 \leq x \leq 3 \\ mx + 2 & , 3 < x \leq 5 \end{cases}$ is differentiable, then the value of $k + m$ is
 (a) 2 (b) $\frac{16}{5}$
 (c) $\frac{10}{3}$ (d) 4

12. The normal to the curve $x^2 + 2xy - 3y^2 = 0$ at $(1, 1)$
 (a) does not meet the curve again
 (b) meets the curve again in the second quadrant
 (c) meets the curve again in the third quadrant
 (d) meets the curve again in the fourth quadrant

13. Let $f(x)$ be a polynomial of degree four having extreme values at $x = 1$ and $x = 2$. If $\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{x^2} \right] = 3$, then $f(2)$ is equal to
 (a) -8 (b) -4
 (c) 0 (d) 4

14. The integral $\int \frac{dx}{x^2(x^4 + 1)^{\frac{3}{4}}}$ equals
 (a) $\left(\frac{x^4 + 1}{x^4} \right)^{\frac{1}{4}} + c$ (b) $(x^4 + 1)^{\frac{1}{4}} + c$
 (c) $-(x^4 + 1)^{\frac{1}{4}} + c$ (d) $-\left(\frac{x^4 + 1}{x^4} \right)^{\frac{1}{4}} + c$

15. The integral $\int_2^4 \frac{\log x^2}{2 \log x^2 + \log(36 - 12x + x^2)} dx$ is equal to
 (a) 2 (b) 4
 (c) 1 (d) 6

16. The area (in sq units) of the region described by $\{(x, y) : y^2 \leq 2x \text{ and } y \geq 4x - 1\}$ is
 (a) $\frac{7}{32}$ (b) $\frac{5}{64}$ (c) $\frac{15}{64}$ (d) $\frac{9}{32}$

17. Let $y(x)$ be the solution of the differential equation $(x \log x) \frac{dy}{dx} + y = 2x \log x$, ($x \geq 1$). Then, $y(e)$ is equal to
 (a) e (b) 0
 (c) 2 (d) $2e$

18. The number of points having both coordinates as integers that lie in the interior of the triangle with vertices $(0, 0)$, $(0, 41)$ and $(41, 0)$ is
 (a) 901 (b) 861
 (c) 820 (d) 780

10 JEE Main Solved Paper 2015

19. Locus of the image of the point (2, 3) in the line $(2x - 3y + 4) + k(x - 2y + 3) = 0$, $k \in \mathbb{R}$, is a
- straight line parallel to X-axis.
 - straight line parallel to Y-axis.
 - circle of radius $\sqrt{2}$.
 - circle of radius $\sqrt{3}$.
20. The number of common tangents to the circles $x^2 + y^2 - 4x - 6y - 12 = 0$ and $x^2 + y^2 + 6x + 18y + 26 = 0$ is
- 1
 - 2
 - 3
 - 4
21. The area (in sq units) of the quadrilateral formed by the tangents at the end points of the latera recta to the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$ is
- $\frac{27}{4}$
 - 18
 - $\frac{27}{2}$
 - 27
22. Let O be the vertex and Q be any point on the parabola $x^2 = 8y$. If the point P divides the line segment OQ internally in the ratio 1 : 3, then the locus of P is
- $x^2 = y$
 - $y^2 = x$
 - $y^2 = 2x$
 - $x^2 = 2y$
23. The distance of the point (1, 0, 2) from the point of intersection of the line $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12}$ and the plane $x - y + z = 16$ is
- $2\sqrt{14}$
 - 8
 - $3\sqrt{21}$
 - 13
24. The equation of the plane containing the line $2x - 5y + z = 3$, $x + y + 4z = 5$ and parallel to the plane $x + 3y + 6z = 1$ is
- $2x + 6y + 12z = 13$
 - $x + 3y + 6z = -7$
 - $x + 3y + 6z = 7$
 - $2x + 6y + 12z = -13$
25. Let a , b and c be three non-zero vectors such that no two of them are collinear and $(a \times b) \times c = \frac{1}{3}|b||c|a$. If θ is the angle between vectors b and c , then a value of $\sin \theta$ is
- $\frac{2\sqrt{2}}{3}$
 - $\frac{-\sqrt{2}}{3}$
 - $\frac{2}{3}$
 - $\frac{-2\sqrt{3}}{3}$
26. If 12 identical balls are to be placed in 3 identical boxes, then the probability that one of the boxes contains exactly 3 balls, is
- $\frac{55}{3} \left(\frac{2}{3}\right)^{11}$
 - $55 \left(\frac{2}{3}\right)^{10}$
 - $220 \left(\frac{1}{3}\right)^{12}$
 - $22 \left(\frac{1}{3}\right)^{11}$
27. The mean of the data set comprising of 16 observations is 16. If one of the observation valued 16 is deleted and three new observations valued 3, 4 and 5 are added to the data, then the mean of the resultant data is
- 16.8
 - 16.0
 - 15.8
 - 14.0
28. If the angles of elevation of the top of a tower from three collinear points A , B and C on a line leading to the foot of the tower are 30° , 45° and 60° respectively, then the ratio $AB : BC$ is
- $\sqrt{3} : 1$
 - $\sqrt{3} : \sqrt{2}$
 - $1 : \sqrt{3}$
 - $2 : 3$
29. Let $\tan^{-1} y = \tan^{-1} x + \tan^{-1} \left(\frac{2x}{1-x^2} \right)$, where $|x| < \frac{1}{\sqrt{3}}$. Then, a value of y is
- $\frac{3x - x^3}{1 - 3x^2}$
 - $\frac{3x + x^3}{1 - 3x^2}$
 - $\frac{3x - x^3}{1 + 3x^2}$
 - $\frac{3x + x^3}{1 + 3x^2}$
30. The negation of $\sim s \vee (\sim r \wedge s)$ is equivalent to
- $s \wedge \sim r$
 - $s \wedge (r \wedge \sim s)$
 - $s \vee (r \vee \sim s)$
 - $s \wedge r$