

WITH GRAPH PAPER

केन्द्रीय माध्यमिक शिक्षा बोर्ड, दिल्ली
मानवियर स्कूल सर्टिफिकेट परीक्षा (कक्षा दशमवीं)
परीक्षार्थी प्रवेश-पत्र के अनुसार स्त्र

Mathematics

041

Wednesday 18.03.2015

English

Code Number

65/2/C

Set Number

Ⓐ ● Ⓑ Ⓒ

B D H S C A

संकेत: B = Back (पृष्ठ), H = Handicapped (असक्षम), S = Special (विशेष), C = Candidate (उम्मीदवार), A = Answer (उत्तर)

संकेत: D = Deaf (श्रवणहीन), H = Hearing Impaired (श्रवण क्षीण), S = Physically Challenged (शारीरिक रूप से असक्षम), C = Candidate (उम्मीदवार), A = Answer (उत्तर)

संकेत: H = Handicapped (असक्षम) संकेत: Yes/No

संकेत: H = Handicapped (असक्षम) संकेत: Yes/No

प्रत्येक अक्षर को एक अक्षर लिखें। नाम के प्रत्येक अक्षर से बीच एक खाना छोड़ दें। यदि सीधे नाम लिखें तो प्रत्येक अक्षर को एक अक्षर लिखें।
Each letter be written in one box and one box be left blank between each part of the name. In case Candidate's Name exceeds 24 letters, write first 24 letters.

041-0852 4474874

य
2A1
7
4
0
14
5
24
6
34
—
—
17
18
19
20
25
26
27

100 - C

100 - C

100 - C

Linear differential Equations

100 - C

Section - C

Q: 20

$$(1+x^2) \frac{dy}{dx} = e^{m \tan^{-1} x} - y$$

$$(1+x^2) \frac{dy}{dx} + y = e^{m \tan^{-1} x}$$

$$\frac{dy}{dx} + \frac{y}{1+x^2} = \frac{e^{m \tan^{-1} x}}{1+x^2}$$

comparing it with

$$\frac{dy}{dx} + Py = Q \rightarrow \text{linear differential equation}$$

$$P = \frac{1}{1+x^2}$$

$$\begin{aligned} \text{I.F.} &= e^{\int P dx} \\ &= e^{\int \frac{1}{1+x^2} dx} \end{aligned}$$

$$\text{I.F.} = e^{\tan^{-1} x}$$

$$\left\{ \int \frac{1}{x^2+1} dx = \tan^{-1} x \right.$$

So the equation becomes

$$y \text{ I.F} = \int Q \times \text{I.F} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{\tan^{-1}x}}{1+x^2} \times e^{\tan^{-1}x} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{(m+1)\tan^{-1}x}}{1+x^2} \, dx + c$$

$$\text{put } \tan^{-1}x = t$$

$$\frac{1}{1+x^2} \, dx = dt$$

$$y e^{\tan^{-1}x} = \int e^{(m+1)t} x \, dt + c$$

$$y e^{\tan^{-1}x} = \frac{e^{(m+1)t}}{(m+1)} + c$$

$$\text{So } y e^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{m+1} + c$$

$$\text{when } x=0 \quad y=1$$

$$ye^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{(m+1)} + c$$

when $x=0$ $y=1$

$$1 \times e^{\tan^{-1}0} = \frac{e^{(m+1)\tan^{-1}0}}{(m+1)} + c$$

$$1 \times e^0 = \frac{e^{(m+1)0}}{m+1} + c \quad e^0 = 1$$

$$1 = \frac{1}{m+1} + c$$

$$1 - \frac{1}{m+1} = c$$

$$\frac{m+1-1}{m+1} = c$$

$$c = \frac{m}{m+1}$$

So equation is

$$\left[ye^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{(1+m)} + \frac{m}{m+1} \right]$$

Q: 21.

$$f(x) = \sin^2 x - \cos x \quad x \in [0, \pi]$$

$$f'(x) = 2 \sin x \cos x - (-\sin x)$$

$$f'(x) = 2 \sin x \cos x + \sin x$$

$$\text{put } f'(x) = 0$$

$$\text{then } (2 \sin x \cos x + \sin x) = 0$$

$$\sin x (2 \cos x + 1) = 0$$

$$\sin x = 0 \quad \text{or} \quad \cos x = -\frac{1}{2}$$

$$x = 0, \pi \quad \text{or} \quad x = \frac{2\pi}{3}$$

$$\text{so } f(0) = [\sin(0)]^2 - \cos 0$$

$$= 0 - 1$$

$$= -1$$

$$f(\pi) = \sin^2 \pi - \cos \pi$$

$$= 0 - (-1)$$

$$= 1$$

$$f\left(\frac{2\pi}{3}\right) = \sin^2 \frac{2\pi}{3} - \cos \frac{2\pi}{3}$$

$$= \left(\frac{\sqrt{3}}{2}\right)^2 - \left(-\frac{1}{2}\right)$$

$$= \frac{3}{4} + \frac{1}{2} = \frac{10}{8} = \frac{5}{4}$$

both the extreme value are automatically included

$$f(0) = -1$$

$$f(\pi) = 1$$

$$f\left(\frac{2\pi}{3}\right) = \frac{5}{4}$$

So Absolute maxima is $\frac{5}{4}$ at $x = \frac{2\pi}{3}$

Absolute minima is -1 at $x = 0$

Q: 22

$$r = \hat{i} + \hat{j} + \hat{k} + \lambda(\hat{i} - \hat{j} + \hat{k})$$

$$r = 4\hat{j} + 2\hat{k} + \mu(2\hat{i} - \hat{j} + 3\hat{k})$$

These lines are coplanar if they are parallel or they are intersecting

But these lines are not parallel.

So they are coplanar if shortest distance between them is zero.

$$\begin{aligned}
 a_1 &= (1, 1, 1) & (a_2 - a_1) &= (-1, 3, 1) \\
 a_2 &= (0, 4, 2) \\
 b_1 &= \langle 1, -1, 1 \rangle \\
 b_2 &= \langle 2, -1, 3 \rangle
 \end{aligned}$$

$$\text{Shortest distance} = \frac{|(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$$

$$\text{let us find } |(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)|$$

$$(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = \begin{vmatrix} -1 & 3 & 1 \\ 1 & -1 & 1 \\ 2 & -1 & 3 \end{vmatrix}$$

(expanding by 1st row)

$$= -1(-3+1) - 3(3-2) + 1(-1+2)$$

$$= -1(-2) - 3(1) + 1$$

$$= 2 + 1 - 3$$

$$= 0$$

So shortest distance between the lines is zero
So they are coplanar.

Let $\langle a, b, c \rangle$ be the direction ratio of normal to the plane.

So the dot product of direction ratio of normal to plane and the direction ratio of line is 0.

$$\begin{aligned} \text{So } a - b + c &= 0 \\ 2a - b + 3c &= 0 \end{aligned}$$

$$\begin{array}{cccc} \underline{a} & \underline{b} & \underline{c} & \\ -1 & 1 & 1 & -1 \\ -1 & 3 & 2 & -1 \end{array}$$

$$\frac{a}{-3+1} = \frac{b}{2-3} = \frac{c}{-1+2}$$

$$\text{So } \langle a, b, c \rangle = \langle -2, -1, 1 \rangle$$

So passing point of line is also the passing point of plane

$$\text{So passing point} = (1, 1, 1)$$

So equation becomes

$$-2(x-1) - 1(y-1) + 1(z-1) = 0$$

$$-2x + 2 - y + 1 + z - 1 = 0$$

$$\boxed{-2x - y + z + 2 = 0}$$

equation of plane.

Q: 23

$$f: W \rightarrow W$$

$$f(n) = \begin{cases} n-1 & \text{if } n \text{ is odd} \\ n+1 & \text{if } n \text{ is even} \end{cases}$$

prove that it is one-one:

let us suppose that $n_1, n_2 \in W$

$$n_1 \neq n_2 \text{ but } f(n_1) = f(n_2)$$

(i) 1st case: if n_1 & n_2 are odd

$$f(n_1) = f(n_2)$$

$$n_1 - 1 = n_2 - 1$$

$$n_1 = n_2$$

which is contradiction.

(ii) 2nd case : n_1, n_2 are even

$$f(n_1) = f(n_2)$$

$$n_1 + 1 = n_2 + 1$$

$$n_1 = n_2$$

which is contradiction.

(iii) 3rd case if n_1 is even & n_2 is odd

$$f(n_1) = f(n_2)$$

$$n_1 + 1 = n_2 - 1$$

$$n_1 + 2 = n_2$$

↓
even

↓
odd

if we add 2 in an even noⁿ then we get an even noⁿ but n_2 is odd which is again contradiction

So from these 3 point we see that

$f(n_1) = f(n_2)$ only if $n_1 = n_2$
 $\Rightarrow f$ is one-one.

To prove that f is onto:

let $y \in W$

and y is even

then $y+1$ is odd and it belongs in whole no.

$$f(y+1) = y+1-1 = y$$

So for every $y \in W$ which is even there exist a preimage $y+1$ which is odd and $(y+1) \in W$

let $y \in W$

y is odd

then $y-1$ is even and $y-1 \in W$

$$f(y-1) = y-1+1 = y$$

So for every $y \in W$ which is odd there exist a pre image $y-1 \in W$ which is even.

So for all $y \in W$ there exist pre image in whole no.
So f is onto.

f is one-one and onto so it is invertible

To find $f^{-1}(x)$

$$y = x - 1 \quad \text{if } x \text{ is odd}$$

$$y + 1 = x$$

y here is even

so $x = y + 1$ if y is even

$$f^{-1}(x) = x + 1 \quad \text{if } x \text{ is even}$$

$$y = x + 1 \quad \text{if } x \text{ is even}$$


y is odd here

$$y - 1 = x$$

so $f^{-1}(x) = x - 1$ if x is odd

$$\text{so } f^{-1}(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$$

$$\boxed{x \in \mathbb{W}} \quad \text{so } \boxed{f^{-1} = f}$$

240

$$y = x - 1 \quad \text{if } x \geq 1$$

$$y = -x + 1 \quad \text{if } x < 1$$

intersection point :

$$y = \sqrt{5-x^2}$$

$$y = x - 1$$

$$\text{so } \sqrt{5-x^2} = (x-1)$$

$$5-x^2 = x^2 + 1 - 2x$$

$$2x^2 - 2x - 4 = 0$$

$$x^2 - x - 2 = 0$$

$$x^2 - 2x + x - 2 = 0$$

$$x(x-2) + 1(x-2) = 0$$

$$\text{so } x = 2 \quad \text{or } x = -1$$

$$x = 2$$

↓ rejected as $x \geq 1$

$$y = \sqrt{5-x^2}$$

$$y = -x+1$$

$$y \quad x < 1$$

$$5-x^2 = x^2+1-2x$$

$$\text{So } x = 2, -1$$

$$x = 2$$

↓
Rejected

$$\boxed{x = -1}$$

Required Area

$$(A) = \int_{-1}^2 \sqrt{5-x^2} dx - \left[\int_{-1}^1 (-x+1) dx + \int_1^2 (x-1) dx \right]$$

$$A = \left. \frac{x\sqrt{5-x^2}}{2} + \frac{5 \sin^{-1} x}{\sqrt{5}} \right|_{-1}^2 - \left[\left. \frac{-x^2+x}{2} \right|_{-1}^1 + \left. \frac{x^2-x}{2} \right|_1^2 \right]$$

$$A = 1 \times 1 + \frac{5 \sin^{-1} 2}{\sqrt{5}} - \left(\frac{-1 \times 2}{2} + \frac{5 \sin^{-1}(-1)}{\sqrt{5}} \right) - \left[\frac{-1}{2} [1-1] + 2 \right. \\ \left. + \frac{1}{2} [4-1] - 1 \right]$$

$$A = 1 + 5 \frac{\sin^{-1} 2}{2} + 1 + 5 \frac{\sin^{-1} 1}{\sqrt{5}} - \left[0 + 2 + \frac{3}{2} - 1 \right]$$

$$A = 2 + 5 \left[\frac{\sin^{-1} 2}{2} + \frac{\sin^{-1} 1}{\sqrt{5}} \right] - \frac{5}{2}$$

$$A = \frac{5 \times \pi}{2} + 2 - \frac{5}{2}$$

$$A = \frac{5\pi}{4} - \frac{1}{2} \text{ sq units}$$

$$\left\{ \begin{aligned} & \frac{\sin^{-1} 2}{\sqrt{5}} + \frac{\sin^{-1} 1}{\sqrt{5}} \\ &= \frac{\sin^{-1} 2 + \sin^{-1} 1}{\sqrt{5}} \\ &= \frac{\pi}{2} \end{aligned} \right\}$$

$\int_1^2 dx$

25°

1st 6 positive integers = (1, 2, 3, 4, 5, 6)

Random Variable = X = larger of 2 numbers.

$\left. \begin{array}{l} x \\ 1 \end{array} \right\}$

Sample space $S = \left\{ \begin{array}{l} (1, 2) (1, 3) (1, 4) (1, 5) (1, 6) \\ (2, 1) (3, 1) (4, 1) (5, 1) (6, 1) \\ (2, 3) (2, 4) (2, 5) (2, 6) (3, 2) (4, 2) (5, 2) \\ (6, 2) (3, 4) (3, 5) (3, 6) (4, 3) (5, 3) (6, 3) \\ (4, 5) (4, 6), (5, 4), (6, 4) (5, 6) (6, 5) \end{array} \right\}$

2

$\{4-1\}-1\}$

X	P(X)	$X_i P_i$	$X_i^2 P_i$	Rough
2	$\frac{2}{30} = \frac{1}{15}$	$\frac{2}{15}$	$\frac{4}{15}$	2+6
3	$\frac{4}{30} = \frac{2}{15}$	$\frac{6}{15}$	$\frac{18}{15}$	
4	$\frac{6}{30} = \frac{3}{15}$	$\frac{12}{15}$	$\frac{48}{15}$	$\frac{236}{15}$
5	$\frac{8}{30} = \frac{4}{15}$	$\frac{20}{15}$	$\frac{100}{15}$	$\frac{48}{15}$
6	$\frac{10}{30} = \frac{5}{15}$	$\frac{30}{15}$	$\frac{180}{15}$	$\frac{200}{15}$

$$\sum X_i P_i = \frac{70}{15} \quad \sum X_i^2 P_i = \frac{350}{15}$$

$$\text{Mean} = \frac{\sum X_i P_i^2}{3} = \frac{14}{3} = 4.66$$

$$\begin{aligned} \text{Variance} &= \sum X_i^2 P_i - (\mu)^2 \\ &= \frac{350}{15} - \frac{70 \times 70}{15} \end{aligned}$$


$$\begin{aligned} \sigma^2 &= \frac{70}{15} \left[\frac{5 - 70}{15} \right] \\ &= \frac{70}{15} \left[\frac{75 - 70}{15} \right] \\ &= \frac{70 \times 5}{3 \times 15 \times 15} \\ &= \frac{14}{9} \end{aligned}$$

$$\sigma^2 = 1.55$$

Variance = 1.55

Section - B

Q:19



$$\begin{array}{ccc} x^x + x^y + y^x = a^b \\ \downarrow \quad \downarrow \quad \downarrow \\ P \quad Q \quad R \end{array}$$

$$\begin{aligned} \frac{d}{dx} [P + Q + R] &= 0 \\ \left[\frac{dP}{dx} + \frac{dQ}{dx} + \frac{dR}{dx} \right] &= 0 \end{aligned}$$

$$P = x^x$$

$$\log P = x \log x$$

$$\frac{1}{P} \frac{dP}{dx} = x \left[\frac{1}{x} \right] + \log x$$

$$\frac{dP}{dx} = x^x (1 + \log x) \quad \text{--- (1)}$$

$$Q = x^y$$

$$\log Q = y \log x$$

$$\frac{1}{Q} \frac{dQ}{dx} = \left(\frac{y}{x} + \log x \frac{dy}{dx} \right)$$

$$\frac{dQ}{dx} = x^y \left[\frac{y}{x} + \log x \frac{dy}{dx} \right] \quad \text{--- (2)}$$

$$Q = y^x$$

$$R = y^x$$

$$\log R = x \log y$$

$$\frac{1}{R} \frac{dR}{dx} = \frac{x}{y} \frac{dy}{dx} + \log y$$

$$\frac{dR}{dx} = y^x \left[\frac{x}{y} \frac{dy}{dx} + \log y \right] \quad \text{--- (3)}$$

adding (1) & (2) & (3) we get

$$x^x + x^x \log x + x^{y-1} y + x^y \log x \frac{dy}{dx} + y^{x-1} x \frac{dy}{dx} + y^x \log y = 0$$

$$\frac{dy}{dx} = \frac{-[x^x(1+\log x) + x^{y-1}y + y^x \log y]}{x^y \log x + y^{x-1}x}$$

Q:18

$$y = e^{ax} \cos bx$$

$$\frac{dy}{dx} = e^{ax} (-\sin bx) b + \cos bx \times e^{ax} \times a$$

$$\frac{dy}{dx} = -b e^{ax} \sin bx + a y \Rightarrow \left[\frac{1}{b} \left(\frac{dy}{dx} - ay \right) = -e^{ax} \sin bx \right] \quad \text{--- (1)}$$

$$\frac{d^2y}{dx^2} = -b [e^{ax} \cos bx \times b + \sin bx \times e^{ax} \times a] + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} = -b \left[by - \frac{a}{b} \left(\frac{dy}{dx} - ay \right) \right] + a \frac{dy}{dx} \quad \text{from (1)}$$

$$\frac{d^2y}{dx^2} = -b^2y + a \frac{dy}{dx} - a^2y + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} + (a^2 + b^2)y - 2a \frac{dy}{dx} = 0$$

hence proved.

26°

$$Z = 5x + 2y$$

↓
objective function to be made maximum and minimum

$$x - 2y \leq 2$$

$$3x + 2y \leq 12$$

$$-3x + 2y \leq 3$$

$$x \geq 0, y \geq 0$$

corresponding equations

$$x - 2y = 2$$

put $(0, 0)$ in it

$$0 - 0 \leq 2$$

which is true so the region is towards

origin

$$3x + 2y \leq 12 \Rightarrow 3x + 2y = 12$$

put $(0, 0)$

$$0 \leq 12 \text{ which is true}$$

so region towards origin

$$-3x + 2y = 3$$

put $(0, 0)$

$$0 \leq 3$$

which is true

so region is towards origin

$x \geq 0$ $y \geq 0$. so it means 1st quadrant.

at A (2,

at B

at C

at

intersection point of

$$-3x + 2y = 3$$

$$3x + 2y = 12$$

$$4y = 15$$

$$y = \frac{15}{4}$$

$$-3x + 2 \times \frac{15}{4} = 3$$

$$-3x = 3 - \frac{15}{2}$$

$$-x = 1 - \frac{5}{2}$$

$$-x = -\frac{3}{2}$$

$$x = \frac{3}{2}$$

$$Z = 5x + 2y$$

at A (2, 0) $Z = 10$

at B (0, 0) $Z = 0$

at C $(0, \frac{3}{2})$ $Z = 3$

at D $(\frac{3}{2}, \frac{15}{4})$ $Z = \frac{15}{2} + \frac{15}{2} = 15$

intersection point

of:

$$x - 2y = 2$$

$$3x + 2y = 12$$

$$4x = 14$$

$$x = \frac{7}{2}$$

$$\frac{7}{2} - 2y = 2$$

$$\frac{7}{2} - 2 = 2y$$

$$\frac{3}{2} = 2y$$

$$y = \frac{3}{4}$$


at E $(\frac{3}{2}, \frac{3}{4})$

$$Z = \frac{15}{2} + \frac{3}{2}$$

$$Z = 9$$

Q126

10 small box = 1 unit



So Z is maximum
at $D(3, 15)$
 $(2, 4)$

$$Z = 15$$

Z is minimum at $(0, 0)$

$$Z = 0$$

170

	Jans	Mats	Toys	
$A =$	30	12	70	→ School X
	40	15	55	→ School Y
	35	20	75	→ School Z
	3×3			

$B =$	25	→ cost of Jans
	100	→ cost of Mats
	50	→ cost of Toys

$$AB = \begin{bmatrix} 30 & 12 & 70 \\ 40 & 15 & 55 \\ 30 & 20 & 75 \end{bmatrix} \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix}$$

$$X = AB = \begin{bmatrix} 750 + 1200 + 3500 \\ 1000 + 1500 + 2750 \\ 875 + 2000 + 3750 \end{bmatrix}$$

$$X = \begin{bmatrix} 5450 \\ 5250 \\ 6625 \end{bmatrix} \begin{array}{l} \rightarrow \text{fund collected by school X} \\ \rightarrow \text{fund collected by school Y} \\ \rightarrow \text{fund collected by school Z} \end{array}$$

fund by X = Rs. 5450

fund by Y = Rs. 5250

fund by Z = Rs. 6625

Total fund = Rs. 17325

They are helping victims and hence

the value of helping others is generated.

$$\underline{16^0} \quad I = \int \frac{x+3}{(x+5)^3} e^x dx$$

$$I = \int \frac{x+5}{(x+5)^3} e^x dx - \int \frac{2 \cdot e^x}{(x+5)^3} dx$$

$$I = \int \frac{1}{(x+5)^2} e^x dx - \int \frac{2 e^x}{(x+5)^3} dx$$

↓
integrating this by parts
 $\frac{1}{(x+5)^2}$ is taken as 1st function

$$I = \frac{1}{(x+5)^2} e^x - \int \frac{d}{dx} \frac{1}{(x+5)^2} e^x dx - \int \frac{2 e^x}{(x+5)^3} dx$$

$$I = \frac{1}{(x+5)^2} e^x + \frac{2}{(x+5)^3} e^x - \int \frac{2 e^x}{(x+5)^3} dx$$

$$\text{So } I = \frac{e^x}{(x+5)^2} + C$$

15°

$$x = a \sin 2t (1 + \cos 2t)$$

$$\frac{dx}{dt} = a [(\sin 2t)(-\sin 2t)(2) + (1 + \cos 2t) \cos 2t \times 2]$$

$$\frac{dx}{dt} = 2a [\cos^2 2t + \cos 2t - \sin^2 2t]$$

$$\frac{dx}{dt} = 2a [\cos 4t + \cos 2t]$$

$$[\cos^2 x - \sin^2 x = \cos 2x]$$

$$y = b \cos 2t (1 - \cos 2t)$$

$$y = b \cos 2t - b \cos^2 2t$$

$$\frac{dy}{dt} = -b \sin 2t \times 2 + b \times 2 \cos 2t \sin 2t \times 2$$

$$\frac{dy}{dt} = 2b [-\sin 2t + \sin 4t]$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dx}{dt}$$

$$\frac{dy}{dx} = \frac{2b}{2a} \left[\frac{\sin 4t - \sin 2t}{\cos 2t + \cos 4t} \right]$$

$$\text{at } t = \frac{\pi}{4}$$

$$\frac{dy}{dx} = \frac{b}{a} \left[\frac{\sin \pi - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2} + \cos \pi} \right]$$

$$= \frac{b}{a} \left[\frac{0 - 1}{0 - 1} \right]$$

$$\left. \frac{dy}{dx} \right|_{t = \frac{\pi}{4}} = \frac{b}{a}$$

140

$$I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2 x}{\sin x + \cos x} dx \quad \text{--- (1)}$$

$$\left[\int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$I = \int_0^{\frac{\pi}{2}} \frac{\sin^2(\frac{\pi}{2}-x)}{\sin(\frac{\pi}{2}-x) + \cos(\frac{\pi}{2}-x)} dx$$

$$I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{\cos x + \sin x} dx \quad \text{--- (2)}$$

adding (1) & (2) we get

$$2I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{\sin^2 x + \cos^2 x}{\sin x + \cos x} dx$$

$$2I = \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{1}{\sin x + \cos x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{1 + \tan^2 x}{2 \tan x + 1 - \tan^2 x} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{2 \tan x + 1 - \tan^2 x} dx$$

put $\sin x = \frac{2 \tan x}{1 + \tan^2 x}$
 and $\cos x = \frac{1 - \tan^2 x}{1 + \tan^2 x}$

put $\tan x = t$

$$\frac{\sec^2 x}{2} dx = dt$$

when $x \rightarrow 0$ $t \rightarrow 0$
 when $x \rightarrow \frac{\pi}{4}$ $t \rightarrow 1$

$$2I = \int_0^1 \frac{dt}{2t + 1 - t^2}$$

$$I = \int_0^1 \frac{dt}{2t + 1 - t^2} \Rightarrow I = \int_0^1 \frac{dt}{2(t-1)^2}$$

$$I = \int_0^1 \frac{dt}{(\sqrt{2})^2 - (t-1)^2} = \int \frac{1}{a^2 - x^2} = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} + t - 1}{\sqrt{2} - t + 1} \right| \Bigg|_0^1$$


$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} + 1 - 1}{\sqrt{2} - 1 + 1} \right| - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log 1 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$= \ominus \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

$$I = -\frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right|$$

13°



$$\Delta = \begin{vmatrix} x+2 & x+6 & x-1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$\Delta = \begin{vmatrix} 3x+7 & 3x+7 & 3x+7 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$\Delta = 3x+7 \begin{vmatrix} 1 & 1 & 1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$C_1 \rightarrow C_1 - C_2 \quad C_2 \rightarrow C_2 - C_3$$

$$\Delta = (3x+7) \begin{vmatrix} 0 & 0 & 1 \\ 7 & -3 & x+2 \\ -3 & -4 & x+6 \end{vmatrix}$$

expanding by R_1

$$\Delta = (3x+7) \begin{vmatrix} -28 & 9 \end{vmatrix}$$

$$\Delta = 0$$

$$\Rightarrow 3x+7=0$$

$$x = \frac{-7}{3}$$

120

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = AA = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1+4+4 & 2+2+4 & 2+4+2 \\ 2+2+4 & 4+1+4 & 4+2+2 \\ 2+4+2 & 4+2+2 & 4+4+1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix} + \begin{bmatrix} -4 & -8 & -8 \\ -8 & -4 & -8 \\ -8 & -8 & -4 \end{bmatrix} + \begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = 0$$

∴ $A^2 - 4A - 5I = 0$

Hence proved.

$$A^2 - 4A - 5I = 0$$

pre-multiplying by A^{-1}

$$A^{-1}AA - 4A^{-1}A - 5A^{-1}I = 0$$

$$IA - 4I - 5A^{-1} = 0$$

$$A - 4I - 5A^{-1} = 0$$

$$A - 4I = 5A^{-1}$$

$$A^{-1} = \frac{A - 4I}{5}$$

$$A^{-1} = \frac{1}{5} [A - 4I]$$

$$5A^{-1} = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} * \begin{bmatrix} -4 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{bmatrix}$$

$$5A^{-1} = \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

$$A^{-1}A = I$$

$$IA = A$$

$$\text{So } A^{-1} = \frac{1}{5} \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

11°

$$\sin^{-1}(1-x) - 2\sin^{-1}x = \frac{\pi}{2}$$

$$\sin^{-1}(1-x) = \frac{\pi}{2} + 2\sin^{-1}x$$

$$(1-x) = \sin\left(\frac{\pi}{2} + 2\sin^{-1}x\right)$$

$$(1-x) = \cos(2\sin^{-1}x)$$

$$2\sin^{-1}x = 0$$

$$x = \sin 0$$

$$\cos 2\theta = 1 - 2\sin^2\theta$$

$$\cos 2\theta = 1 - 2x^2$$

$$\Rightarrow 1-x = \cos 2\theta$$

$$1-x = 1-2x^2$$

$$\Rightarrow -2x^2 - x = 0$$

$$x[2x-1] = 0$$

$$x = 0 \text{ or } x = \frac{1}{2}$$

put $x = \frac{1}{2}$ in equation

$$x \sin^{-1} \frac{1}{2} - 2x \sin^{-1} \frac{1}{2}$$

$$= \frac{1}{6} - 2 \times \frac{1}{6}$$

$$\neq \frac{1}{2}$$

$$\text{So } x \neq \frac{1}{2}$$

$$\text{So } \boxed{x=0}$$

10°

passing point of line = (4, 2, 2)

since \vec{b} is \perp to line

So direction ratio of line = $\langle 2, 3, 6 \rangle$

So equation of line

$$\frac{x-4}{2} = \frac{y-2}{3} = \frac{z-2}{6} = \lambda$$

general point on line

$$(2\lambda+4, 3\lambda+2, 6\lambda+2)$$

$$P(1, 2, 3)$$

Q

line

Q be the foot of \perp ar

PQ is \perp ar to line

So dot product of direction ratios of line and PQ is 0

$$\text{direction ratios of } Q = \langle 2\lambda+3, 3\lambda, 6\lambda-1 \rangle$$

So according to question:

$$2(2\lambda+3) + (3\lambda)3 + 6(6\lambda-1) = 0$$

$$4\lambda + 6 + 9\lambda + 36\lambda - 6 = 0$$

$$\boxed{\lambda = 0}$$

So point $Q = (4, 2, 2)$

So Lar distance

$$PQ = \sqrt{(4-1)^2 + (2-2)^2 + (2-3)^2}$$

$$= \sqrt{(3)^2 + (0)^2 + (-1)^2}$$

$$= \sqrt{9+1}$$

$$\boxed{\text{length of Lar} = \sqrt{10} \text{ unit}}$$

90ABthese A, B, C, D are coplaner :

so $\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = 0$
 triple product is 0.

$$\vec{AB} = 1\hat{i} + (\lambda - 1)\hat{j} + 4\hat{k}$$

$$\vec{BC} = 0\hat{i} + (1 - \lambda)\hat{j} - 7\hat{k}$$

$$\vec{CD} = 2\hat{i} + 3\hat{j} + \hat{k}$$

$$\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = \begin{vmatrix} 1 & \lambda - 1 & 4 \\ 0 & 1 - \lambda & -7 \\ 2 & 3 & 1 \end{vmatrix} = 0$$

expanding by R_1

$$1(1 - \lambda + 21) - (\lambda - 1)14 + 4(2(\lambda - 1)) = 0$$

$$22 - \lambda - 14\lambda + 14 + 8\lambda - 8 = 0 \quad \boxed{\lambda = 4}$$

$$-7\lambda = -28$$

8°

P (probability of success) = $\frac{1}{2}$
i.e. that head comes

Q (probability of failure) = $\frac{1}{2}$
i.e. that tail comes

Let the coin be tossed n times

this event follow the conditions of Bernoulli trial

X be the random variable = no. of heads

$$P(X \geq 1) = P(1) + \dots + P(n)$$

$$P(X \geq 1) = 1 - P(0)$$

$$= 1 - {}^n C_0 \times \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^n$$

$P(X \geq 1)$ should be more than 80%.

$$P(X \geq 1) > \frac{8}{10} \quad \frac{80}{100} < 1 - \left(\frac{1}{2}\right)^n$$

$$1 - \left(\frac{1}{2}\right)^n \geq \frac{8}{10}$$

$$\left(\frac{1-8}{10}\right) \geq \left(\frac{1}{2}\right)^n$$

$$\frac{18}{50} > \left(\frac{1}{2}\right)^n$$

$$5 < 2^n$$

$$n = 3$$

follow the condition
So the coin should be tossed
at least 3 times.

7.

$$I = \int \frac{x^2}{x^4 + x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^4 + 2x^2 - x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^2(x^2 + 2) - 1(x^2 + 2)} dx$$

$$I = \int \frac{x^2}{(x^2-1)(x^2+2)} dx$$

put $x^2 = t$

then
$$\frac{t}{(t-1)(t+2)} = \frac{A}{t-1} + \frac{B}{t+2}$$

$$t = A(t+2) + B(t-1)$$

put $t=1$

$$1 = A \times 3$$

$$\boxed{A = \frac{1}{3}}$$

put $t=-2$

$$-2 = -3B$$

$$\boxed{B = \frac{2}{3}}$$

$$I = \int \frac{1}{3(t-1)} dx + \frac{2}{3} \int \frac{dx}{t+2}$$

$$I = \int \frac{1}{3(x^2-1)} dx + \frac{2}{3} \int \frac{dx}{(x^2+2)} \quad \int \frac{1}{x^2-a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right|$$

$$I = \frac{1}{3} \times \frac{1}{2} \log \left| \frac{x-1}{x+1} \right| + \frac{2}{3} \times \frac{1}{\sqrt{2}} \tan^{-1} \frac{x}{\sqrt{2}} + C \quad \int \frac{1}{x^2+a^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

$$I = \frac{1}{6} \log \left| \frac{x-1}{x+1} \right| + \frac{\sqrt{2}}{3} \tan^{-1} \frac{x}{\sqrt{2}} + C$$

~~Q~~

Section - A

Q: 6

equation of plane

$$6x - 3y + 2z - 4 = 0$$

$$\text{distance} = \frac{|6 \times 2 - 3 \times 5 + 2(-3) - 4|}{\sqrt{36 + 9 + 4}}$$

$$= \frac{|12 - 15 - 6 - 4|}{\sqrt{49}}$$

$$= \frac{|-13|}{7}$$

$$\boxed{\text{distance} = \frac{13}{7} \text{ units}}$$

Q: 5

$$\vec{a} = \hat{i} - \hat{j}$$

$$|\vec{a}| = \sqrt{2}$$

$$\vec{b} = \hat{j} - \hat{k}$$

$$|\vec{b}| = \sqrt{2}$$

0902

Fictitious Roll No.
(To be entered by Board)

41474874

अपना अनुक्रमीक इस उत्तर-पुस्तिका
पर न लिखें
Please do not write your
Roll Number on this Answer-Book

अतिरिक्त उत्तर-पुस्तिका (सी) की सहायता
Supplementary Answer-Book (Series)

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$$

$$(\hat{i} - \hat{j}) \cdot (\hat{j} - \hat{k}) = \sqrt{2} \times \sqrt{2} \cos \theta$$

$$\frac{-1}{2} = \cos \theta$$

$$\theta = \frac{2\pi}{3}$$

angle between vectors = $\frac{2\pi}{3}$

40

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 3 \\ 3 & 5 & -2 \end{vmatrix}$$

$$\vec{a} \times \vec{b} = \hat{i}(-2-15) - \hat{j}(-4-9) + \hat{k}(10-3)$$

$$\vec{a} \times \vec{b} = -17\hat{i} + 13\hat{j} + 7\hat{k}$$

$$|\vec{a} \times \vec{b}| = \sqrt{(-17)^2 + (13)^2 + (7)^2}$$

$$= \sqrt{289 + 169 + 49}$$

$$= \sqrt{507}$$

$$= \sqrt{3 \times 169}$$

$$|\vec{a} \times \vec{b}| = 13\sqrt{3}$$

Q: 3

$$x \log x \frac{dy}{dx} + y = 2 \log x$$

$$\frac{dy}{dx} + \frac{y}{x \log x} = 2 \log x$$

compare it with $\frac{dy}{dx} + Py = Q$

$$I.F = e^{\int P dx}$$

$$= e^{\int x \log x dx}$$

put $\log x = t$
 $\frac{1}{x} dx = dt$

$$I \cdot F = e^{\int \frac{dt}{t}}$$

$$= e^{\log |t|}$$

$$= t$$

$$\boxed{I \cdot F = \log x}$$

2° general equation of family of lines passing through origin

$$y = mx$$

$$m = \frac{y}{x}$$

$$\frac{dy}{dx} = m$$

$$\frac{dy}{dx} = \frac{y}{x}$$

$$\boxed{x \frac{dy}{dx} - y = 0}$$

Q.10

$$a_{12} = e^{2x} \sin x$$

$$a_{12} = e^{2x} \sin x$$

POANAM 01/07/2019/A
Evaluation done as per CBSE marking scheme.

XII - U